

**THE FIRST COHOMOLOGY GROUP OF THE GENERALIZED  
MORAVA STABILIZER ALGEBRA  
(DRAFT VERSION)**

HIROFUMI NAKAI AND DOUGLAS C. RAVENEL  
September 8, 2000

ABSTRACT. There are  $p$ -local spectra  $T(m)$  with  $BP_*(T(m)) = BP_*[t_1, \dots, t_m]$ . Its Adams-Novikov  $E_2$ -term is isomorphic to

$$\mathrm{Ext}_{\Gamma(m+1)}(BP_*, BP_*),$$

where

$$\Gamma(m+1) = BP_*(BP)/(t_1, \dots, t_m) = BP_*[t_{m+1}, t_{m+2}, \dots].$$

In this paper we determine the groups

$$\mathrm{Ext}_{\Gamma(m+1)}^1(BP_*, v_n^{-1}BP_*/I_n)$$

for all  $m, n > 0$ . Its rank ranges from  $2n+1$  to  $n^2$  depending on the value of  $m$ .

1. INTRODUCTION AND MAIN THEOREM

Let  $BP$  be the Brown-Peterson spectrum for a fixed prime  $p$ . In [Rav86, §6.5], the second author has introduced the spectrum  $T(m)$  which has  $BP_*$ -homology

$$BP_*(T(m)) = BP_*[t_1, \dots, t_m],$$

and is homotopy equivalent to  $BP$  below dimension  $2p^{m+1} - 3$ .

Then the Adams-Novikov  $E_2$ -term converging to the homotopy groups of  $T(m)$

$$E_2^{*,*}(T(m)) = \mathrm{Ext}_{BP_*(BP)}(BP_*, BP_*(T(m)))$$

is isomorphic by [Rav86, 7.1.3] to

$$\mathrm{Ext}_{\Gamma(m+1)}(BP_*, BP_*),$$

where

$$\Gamma(m+1) = BP_*(BP)/(t_1, \dots, t_m) = BP_*[t_{m+1}, t_{m+2}, \dots].$$

In particular  $\Gamma(1) = BP_*(BP)$  by definition. To get the structure of this, we can use the chromatic method introduced in [MRW77].

Recall the Morava stabilizer algebra

$$\Sigma(n) = K(n)_* \otimes_{BP_*} BP_*(BP) \otimes_{BP_*} K(n)_*$$

and the isomorphism ([MR77] and [Rav86, 6.1.1])

$$\mathrm{Ext}_{BP_*(BP)}(BP_*, v_n^{-1}BP_*/I_n) \cong \mathrm{Ext}_{\Sigma(n)}(K(n)_*, K(n)_*).$$

As an algebra,

$$\Sigma(n) = K(n)_*[t_1, t_2, \dots]/(v_n t_i^{p^n} - v_n^{p^i} t_i),$$

---

The second author acknowledges support from NSF grant DMS-9802516.

where  $t_i$  is the image of the generator of the same name in  $BP_*(BP)$ . As in [Rav86, §6.5] we let

$$\Sigma(n, m+1) = \Sigma(n)/(t_1, \dots, t_m);$$

we call this the *generalized Morava stabilizer algebra*. The object of this paper is to determine its first cohomology group,

$$\mathrm{Ext}_{\Sigma(n, m+1)}^1(K(n)_*, K(n)_*)$$

(which we will abbreviate by  $\mathrm{Ext}_{\Sigma(n, m+1)}^1$ ) for all values of  $m \geq 0$  and  $n > 0$  and for all primes  $p$ . This amounts to identifying the primitive elements in  $\Sigma(n, m+1)$ . The case  $m = 0$  was described in [Rav86, 6.3.12].

As explained in [Rav86, §6.2], the cohomology of  $\Sigma(n)$  is essentially the continuous cohomology of a certain pro- $p$ -group  $S_n$  known as the Morava stabilizer group. It can be described as a group of automorphisms of a certain formal group law  $F_n$  in characteristic  $p$  and as a group of units in the maximal order  $E_n$  of a certain  $p$ -adic division algebra  $D_n$ .  $E_n$  is also the endomorphism ring of  $F_n$ .

In a similar way  $\Sigma(n, m+1)$  is related to a subgroup of  $S_n$ . In terms of the formal group law it is the subgroup of automorphisms given by power series congruent to the variable  $x$  modulo  $(x^{p^{m+1}})$ . In terms of  $E_n$  it is the multiplicative group of units congruent to 1 modulo the ideal  $(S^{m+1})$ .

The ring  $E_n$  has an embedding in the ring of  $n \times n$  matrices over the Witt ring  $W(\mathbf{F}_{p^n})$  described in [Rav86, 6.2.6]. This means that  $S_n$  and each of its subgroups supports a homomorphism induced by the determinant to the group of units in  $W(\mathbf{F}_{p^n})$ , and it is known that its image is contained in the  $p$ -adic units  $\mathbf{Z}_p^\times$ . The structure of this group is

$$\mathbf{Z}_p^\times \cong \begin{cases} \mathbf{Z}/(p-1) \oplus \mathbf{Z}_p & \text{for } p \text{ odd} \\ \mathbf{Z}/(2) \oplus \mathbf{Z}_2 & \text{for } p = 2. \end{cases}$$

From this it is possible to construct primitives  $T_n \in \Sigma(n)$  for all primes  $p$  and  $U_n \in \Sigma(n)$  for  $p = 2$  [Rav86, 6.3.12] satisfying

$$\begin{aligned} T_n &\equiv \sum_{0 \leq j < n} t_n^{p^j} \pmod{(t_1, \dots, t_{n-1})} \\ \text{and } U_n - T_n &\equiv \sum_{0 \leq j < n} t_{2n}^{2^j} \pmod{(t_1, \dots, t_{n-1})}. \end{aligned}$$

The corresponding elements in  $\mathrm{Ext}_{\Sigma(n)}^1$ , and their images in  $\mathrm{Ext}_{\Sigma(n, m+1)}^1$ , are denoted by  $\zeta_n$  and  $\rho_n$  respectively.

The results of [Rav86, §6.3] are stated in terms of  $S(n) = \Sigma(n) \otimes_{K(n)_*} \mathbf{F}_p$  and  $S(n, m+1) = \Sigma(n, m+1) \otimes_{K(n)_*} \mathbf{F}_p$ . Passing from  $\Sigma(n)$  to  $S(n)$  amounts to dropping the grading and setting  $v_n$  equal to 1. Formulas are given for  $T_n$  and (for  $p = 2$ )  $U_n$  in  $S(n)$ . It is straightforward to lift them to homogeneous elements in  $\Sigma(n)$ .

We can now state our main result.

**Theorem 1.1.** For  $p$  odd the rank of  $\text{Ext}_{\Sigma(n,m+1)}^1$  (as a vector space over  $K(n)_*$ ) is

$$\begin{cases} (m+1)n+1 & \text{for } m < \frac{n-2}{2} \\ (m+1)n+n/2 & \text{for } n \text{ even and } m = \frac{n-2}{2} \\ (m+1)n & \text{for } \frac{n-1}{2} \leq m \leq n-1 \\ n^2 & \text{for } m \geq n-1. \end{cases}$$

Let  $h_{m+i,j} \in \text{Ext}^1$  be the element corresponding to  $t_{m+i}^{p^j}$  when it is primitive. Then a basis for  $\text{Ext}^1$  is given by

$$\begin{cases} \{\zeta_n\} \cup \{h_{m+i,j} : 1 \leq i \leq m+1, j \in \mathbf{Z}/(n)\} & \text{for } m < \frac{n-2}{2} \\ \begin{cases} \{\zeta_{n,j} : j \in \mathbf{Z}/(n/2)\} \\ \cup \{h_{m+i,j} : 1 \leq i \leq m+1, j \in \mathbf{Z}/(n)\} \end{cases} & \text{for } n \text{ even and } m = \frac{n-2}{2} \\ \{h_{m+i,j} : 1 \leq i \leq m+1, j \in \mathbf{Z}/(n)\} & \text{for } \frac{n-1}{2} \leq m \leq n-1 \\ \{h_{m+i,j} : 1 \leq i \leq n, j \in \mathbf{Z}/(n)\} & \text{for } m \geq n. \end{cases}$$

where  $\zeta_n$  is as above and

$$\zeta_{n,j} = v_n^{-p^j} (t_n + v_n^{1-p^{n/2}} t_n^{p^{n/2}} - t_{n/2}^{1+p^{n/2}})^{p^j}.$$

For  $p=2$  the rank is

$$\begin{cases} (m+1)n+2 & \text{for } m < \frac{n-2}{2} \\ (m+1)n+n/2+1 & \text{for } n \text{ even and } m = \frac{n-2}{2} \\ (m+1)n+1 & \text{for } \frac{n-1}{2} \leq m \leq n-1 \\ n^2 & \text{for } m \geq n. \end{cases}$$

The basis is as in the odd primary case but with  $\rho_n$  added when  $m < n$ .

Note that for  $m=0$  this result gives the same answer as [Rav86, 6.3.12]. Also [Rav86, 6.5.6] implies that  $\text{Ext}^1$  has rank  $n^2$  with the basis indicated above when  $m > \frac{pm}{2p-2} - 1$  and  $m \geq n-1$ ; it says that in that case the full Ext group is the exterior algebra on those generators. [There is a missing hypothesis in [Rav86, 6.5.6] and [Rav86, 6.3.7]; see the online errata for details.]

**Corollary 1.2.** For  $n \leq 3$  the rank of  $\text{Ext}_{\Sigma(n,m+1)}^1$  is as indicated in the following table.

$n=1$				$n=2$				$n=3$			
$p=2$		$p \text{ odd}$		$p=2$		$p \text{ odd}$		$p=2$		$p \text{ odd}$	
$m$	rank	$m$	rank	$m$	rank	$m$	rank	$m$	rank	$m$	rank
0	2	$\geq 0$	1	0	4	0	3	0	5	0	4
$\geq 1$	1			1	5	$\geq 1$	4	1	7	1	6
				$\geq 2$	4			2	10	$\geq 2$	9
								$\geq 3$	9		

## 2. THE PROOF

We need to show that the indicated basis elements are primitive and that there are no other primitives. The primitivity of  $\zeta_n$  and (for  $p = 2$ )  $\rho_n$  was established in [Rav86, 6.3.12].

For the rest we need to study the coproduct in  $\Sigma(n, m + 1)$ . A formula for the coproduct in  $BP_*(BP)$  was given in [Rav86, 4.3.13]. In  $BP_*(BP)/I_n$  for  $i \leq 2n$  we have [Rav86, 4.3.15]

$$\Delta(t_i) = \sum_{0 \leq j \leq i} t_j \otimes t_{i-j}^{p^j} + \sum_{0 \leq j \leq i-n-1} v_{n+j} b_{i-n-j, n+j-1},$$

where  $b_{i,j}$  satisfies

$$b_{i,j} \equiv -\frac{1}{p} \sum_{0 < k < p^{j+1}} \binom{p^{j+1}}{k} t_i^k \otimes t_i^{p^{j+1}-k} \pmod{(t_1, \dots, t_{i-1})}.$$

It is defined precisely in [Rav86, 4.3.14]. Similar methods yield the following formula for the coproduct in  $\Gamma(m + 1)/I_n$  for  $i \leq 2n$ .

$$\begin{aligned} \Delta(t_{m+i}) &= t_{m+i} \otimes 1 + 1 \otimes t_{m+i} + \sum_{m < k < i} t_k \otimes t_{m+i-k}^{p^k} \\ &\quad + \sum_{0 \leq k \leq i-n-1} v_{n+k} b_{m+i-n-k, n+k-1}. \end{aligned}$$

In  $\Sigma(n, m + 1)$  this simplifies to

$$(2.1) \quad \begin{aligned} \Delta(t_{m+i}) &= t_{m+i} \otimes 1 + 1 \otimes t_{m+i} + \sum_{m < k < i} t_k \otimes t_{m+i-k}^{p^k} \\ &\quad + v_n b_{m+i-n, n-1}, \end{aligned}$$

where the last term vanishes when  $i \leq n$ . This formula implies that  $t_{m+i}$  is primitive for  $i \leq \min(m + 1, n)$ .

When  $n$  is even and  $m = \frac{n-2}{2}$  we have

$$\begin{aligned} \Delta(t_n) &= t_n \otimes 1 + 1 \otimes t_n + t_{n/2} \otimes t_{n/2}^{p^{n/2}}, \\ \Delta(v_n^{1-p^{n/2}} t_n^{p^{n/2}}) &= v_n^{1-p^{n/2}} \left( t_n \otimes 1 + 1 \otimes t_n + t_{n/2} \otimes t_{n/2}^{p^{n/2}} \right)^{p^{n/2}} \\ &= v_n^{1-p^{n/2}} \left( t_n^{p^{n/2}} \otimes 1 + 1 \otimes t_n^{p^{n/2}} + t_{n/2}^{p^{n/2}} \otimes t_{n/2}^{p^{n/2}} \right) \\ &= v_n^{1-p^{n/2}} \left( t_n^{p^{n/2}} \otimes 1 + 1 \otimes t_n^{p^{n/2}} + v_n^{p^{n/2}-1} t_{n/2}^{p^{n/2}} \otimes t_{n/2} \right) \\ &= v_n^{1-p^{n/2}} \left( t_n^{p^{n/2}} \otimes 1 + 1 \otimes t_n^{p^{n/2}} \right) + t_{n/2}^{p^{n/2}} \otimes t_{n/2}, \\ \text{and } \Delta(t_{n/2}^{1+p^{n/2}}) &= \left( t_{n/2} \otimes 1 + 1 \otimes t_{n/2} \right)^{1+p^{n/2}} \\ &= t_{n/2}^{1+p^{n/2}} \otimes 1 + t_{n/2}^{p^{n/2}} \otimes t_{n/2} + t_{n/2} \otimes t_{n/2}^{p^{n/2}} + 1 \otimes t_{n/2}^{1+p^{n/2}}, \end{aligned}$$

so  $\zeta_{n,j}$  is primitive.

*This means that each basis element specified in Theorem 1.1 is indeed primitive.*

To show that there are no other primitives in  $\Sigma(n, m + 1)$  we need the methods of [Rav86, §6.3]. As noted above, results there are stated in terms of  $S(n) = \Sigma(n) \otimes_{K(n)_*} \mathbf{F}_p$  and  $S(n, m + 1) = \Sigma(n, m + 1) \otimes_{K(n)_*} \mathbf{F}_p$ . An increasing filtration

on  $S(n)$  is described in [Rav86, 6.3.1]. The weight of  $t_i^{p^j}$  for each  $j$  is the integer  $d_{n,i}$  defined recursively by

$$d_{n,i} = \begin{cases} 0 & \text{if } i \leq 0 \\ \max(i, pd_{n,i-n}) & \text{if } i > 0. \end{cases}$$

The bigraded object  $E^0S(n)$  is described in [Rav86, 6.3.2]. It is considerably simpler than the coproduct in the unfiltered object. It contains elements  $t_{m+i,j}$  (with  $j \in \mathbf{Z}/(n)$ ) which are the projections of  $t_{m+i}^{p^j}$ . The coproduct on these elements is given by

$$(2.2) \quad \Delta(t_{m+i,j}) = \begin{cases} t_{m+i,j} \otimes 1 + 1 \otimes t_{m+i,j} \\ \quad + \sum_{m < k < i} t_{k,j} \otimes t_{m+i-k,j+k} & \text{if } i < c - m \\ t_{m+i,j} \otimes 1 + 1 \otimes t_{m+i,j} \\ \quad + \sum_{m < k < i} t_{k,j} \otimes t_{m+i-k,j+k} \\ \quad + \bar{b}_{m+i-n,n-1+j} & \text{if } i = c - m \\ t_{m+i,j} \otimes 1 + 1 \otimes t_{m+i,j} \\ \quad + \bar{b}_{m+i-n,n-1+j} & \text{if } i > c - m. \end{cases}$$

where  $c = pn/(p-1)$  and  $\bar{b}_{m+i-n,n-1+j}$  is the projection of  $b_{m+i-n,n-1+j}$ , which is 0 for  $i \leq n$ .

Note  $t_{m+i,j}$  is primitive for  $i \leq m+1$  as expected, but it is also primitive for  $c-m < i \leq n$ , which can occur when  $m > n/(p-1)$ .

To proceed further we use the fact that the dual of  $E^0S(n, m+1)$  is a primitively generated Hopf algebra and therefore isomorphic to the universal enveloping algebra of its restricted Lie algebra of primitives, by a theorem of Milnor-Moore [MM65]. The cohomology of the unrestricted Lie algebra  $L(n, m+1)$  (this notation differs from that of [Rav86, §6.3]) is that of the Koszul complex

$$(2.3) \quad C(n, m+1) = E(h_{m+i,j} : i > 0, j \in \mathbf{Z}/(n)),$$

where each  $h_{m+i,j}$  has cohomological degree 1, with

$$d(h_{m+i,j}) = \begin{cases} \sum_{m < k < i} h_{k,j} h_{m+i-k,j+k} & \text{if } i \leq c - m \\ 0 & \text{if } i > c - m. \end{cases}$$

**Lemma 2.4.** *Let  $C(n, m+1)$  be the complex of (2.3). Then  $H^1(L(n, m+1)) = H^1(C(n, m+1))$  is spanned by*

$$\{h_{m+i,j} : 1 \leq i \leq m+1\} \cup \{h_{m+i,j} : i > c - m\} \cup \left\{ \sum_j h_{n,j}, \sum_j h_{2n,j} \right\},$$

(where  $c = pn/(p-1)$ ) unless  $n = 2m+2$ , in which case we must adjoin the set

$$\{h_{n,j} + h_{n,j+n/2} : j \in \mathbf{Z}/(n/2)\}.$$

Note that  $h_{n,j}$  is either trivial or in the first subset unless  $n \geq 2m+2$  and that  $h_{n,j}$  is either trivial or in the second subset unless  $p = 2$ . Note also that the first and second subsets overlap when  $m \geq c/2$ .

*Proof.* The primitivity of the elements in the first and second subsets is obvious. For  $\sum_j h_{n,j}$  we have

$$\begin{aligned}
d\left(\sum_j h_{n,j}\right) &= \sum_j \sum_{m < k < n-m} h_{k,j} h_{n-k,j+k} \\
&= \sum_{m < k < n/2} \sum_j h_{k,j} h_{n-k,j+k} \\
&\quad + \begin{cases} \sum_j h_{n/2,j} h_{n/2,j+n/2} & \text{if } n \text{ is even} \\ 0 & \text{if } n \text{ is odd} \end{cases} \\
&\quad + \sum_{n/2 < k < n-m} \sum_j h_{k,j} h_{n-k,j+k} \\
&= \sum_{m < k < n/2} \sum_j h_{k,j} h_{n-k,j+k} + h_{n-k,j+k} h_{k,j} \\
&\quad + \begin{cases} \sum_{0 \leq j < n/2} h_{n/2,j} h_{n/2,j+n/2} \\ \quad + \sum_{n/2 \leq j < n} h_{n/2,j} h_{n/2,j+n/2} & \text{if } n \text{ is even} \\ 0 & \text{if } n \text{ is odd} \end{cases} \\
&= \begin{cases} \sum_{0 \leq j < n/2} h_{n/2,j} h_{n/2,j+n/2} + h_{n/2,j+n/2} h_{n/2,j} & \text{if } n \text{ is even} \\ 0 & \text{if } n \text{ is odd} \end{cases} \\
&= 0.
\end{aligned}$$

Similar calculations show that for  $p = 2$ ,  $\sum_j h_{2n,j}$  is a cocycle, and that for  $n = 2m + 2$ ,  $h_{n,j} + h_{n,j+n/2}$  is one.

It remains to show that there are no other cocycles in the subspace spanned by

$$\{h_{m+i,j} : m+1 < i \leq c-m\},$$

which is nonempty only when

$$m < \frac{pn - p + 1}{2(p-1)}.$$

It suffices to consider elements which are homogeneous with respect to the filtration grading, i.e., to restrict our attention to one value of  $i$  at a time. Thus we need to show that the subspace spanned by

$$(2.5) \quad \left\{ \sum_{m < k < i} h_{k,j} h_{m+i-k,j+k} : j \in \mathbf{Z}/(n) \right\}$$

has dimension

$$(2.6) \quad \begin{cases} n/2 & \text{if } m+i = n \text{ and } n = 2m+2 \\ n-1 & \text{if } m+i = n \text{ and } n > 2m+2 \\ n-1 & \text{if } m+i = 2n \\ n & \text{otherwise.} \end{cases}$$

When  $n = 2m + 2$  and  $m + i = n$ , the set of (2.5) is

$$\begin{aligned} & \{h_{n/2,j}h_{n/2,j+n/2} : j \in \mathbf{Z}/(n)\} \\ &= \{h_{n/2,j}h_{n/2,j+n/2} : 0 \leq j < n/2\} \cup \{h_{n/2,j}h_{n/2,j+n/2} : n/2 \leq j < n\} \\ &= \{h_{n/2,j}h_{n/2,j+n/2} : 0 \leq j < n/2\} \cup \{-h_{n/2,j+n/2}h_{n/2,j} : n/2 \leq j < n\} \\ &= \{h_{n/2,j}h_{n/2,j+n/2} : 0 \leq j < n/2\} \cup \{-h_{n/2,j}h_{n/2,j+n/2} : 0 \leq j < n/2\}, \end{aligned}$$

so the subspace it spans has dimension  $n/2$ .

Now suppose that  $m + i = n$ ,  $n > 2m + 2$ , and  $n$  is odd. It suffices to consider the middle two terms in the sum. Let  $\ell = (n - 1)/2$ . Then we have

$$d(h_{n,j}) = h_{\ell,j}h_{\ell+1,j+\ell} + h_{\ell+1,j}h_{\ell,j+\ell+1} + \dots$$

We can cancel the second term by adding  $d(h_{n,j+\ell+1})$ , i.e.,

$$\begin{aligned} & d(h_{n,j} + h_{n,j+\ell+1}) \\ &= h_{\ell,j}h_{\ell+1,j+\ell} + h_{\ell+1,j}h_{\ell,j+\ell+1} \\ &\quad + h_{\ell,j+\ell+1}h_{\ell+1,j+\ell+1} + h_{\ell+1,j+\ell+1}h_{\ell,j+\ell+1+\ell+1} + \dots \\ &= h_{\ell,j}h_{\ell+1,j+\ell} + h_{\ell+1,j}h_{\ell,j+\ell+1} \\ &\quad + h_{\ell,j+\ell+1}h_{\ell+1,j} + h_{\ell+1,j+\ell+1}h_{\ell,j+1} + \dots \\ &= h_{\ell,j}h_{\ell+1,j+\ell} + h_{\ell+1,j+\ell+1}h_{\ell,j+1} + \dots \end{aligned}$$

Similarly we can cancel the second term here by adding  $d(h_{n,j+1})$ . Since  $(n + 1)/2$  and  $n$  are relatively prime, we will need to sum up the  $h_{n,j}$  over all  $j$  to get a cocycle. It follows that this subspace has dimensions  $n - 1$  as claimed.

For  $m + i = n$  and  $n$  even, let  $\ell = n/2$ . Then it suffices to consider the middle three terms of the sum, i.e.,

$$d(h_{n,j}) = h_{\ell-1,j}h_{\ell+1,j+\ell-1} + h_{\ell,j}h_{\ell,j+\ell} + h_{\ell+1,j}h_{\ell-1,j+\ell+1} + \dots$$

We can cancel the middle term by adding  $d(h_{n,j+\ell})$ , so we get

$$\begin{aligned} & d(h_{n,j} + h_{n,j+\ell}) \\ &= h_{\ell-1,j}h_{\ell+1,j+\ell-1} + h_{\ell,j}h_{\ell,j+\ell} + h_{\ell+1,j}h_{\ell-1,j+\ell+1} \\ &\quad + h_{\ell-1,j+\ell}h_{\ell+1,j-1} + h_{\ell,j+\ell}h_{\ell,j} + h_{\ell+1,j+\ell}h_{\ell-1,j+1} + \dots \\ &= h_{\ell-1,j}h_{\ell+1,j+\ell-1} + h_{\ell-1,j+\ell}h_{\ell+1,j-1} \\ &\quad + h_{\ell+1,j}h_{\ell-1,j+\ell+1} + h_{\ell+1,j+\ell}h_{\ell-1,j+1} + \dots \end{aligned}$$

Now we can cancel the third and fourth terms by adding  $d(h_{n,j+1} + h_{n,j+\ell+1})$ , and we have

$$\begin{aligned} & d(h_{n,j} + h_{n,j+\ell} + h_{n,j+1} + h_{n,j+\ell+1}) \\ &= h_{\ell-1,j}h_{\ell+1,j+\ell-1} + h_{\ell-1,j+\ell}h_{\ell+1,j-1} \\ &\quad + h_{\ell+1,j}h_{\ell-1,j+\ell+1} + h_{\ell+1,j+\ell}h_{\ell-1,j+1} \\ &\quad + h_{\ell-1,j+1}h_{\ell+1,j+\ell} + h_{\ell-1,j+\ell+1}h_{\ell+1,j} \\ &\quad + h_{\ell+1,j+1}h_{\ell-1,j+\ell+2} + h_{\ell+1,j+\ell+1}h_{\ell-1,j+2} + \dots \\ &= h_{\ell-1,j}h_{\ell+1,j+\ell-1} + h_{\ell-1,j+\ell}h_{\ell+1,j-1} \\ &\quad + h_{\ell+1,j+1}h_{\ell-1,j+\ell+2} + h_{\ell+1,j+\ell+1}h_{\ell-1,j+2} + \dots \end{aligned}$$

Again in order to get complete cancellation we need to sum over all  $j$ , so the subspace has dimension  $n - 1$  as claimed.

We can make a similar argument for  $m + i = 2n$  when  $p = 2$ , namely

$$\begin{aligned} d(h_{2n,j}) &= h_{n-1,j}h_{n+1,j-1} + h_{n,j}h_{n,j} + h_{n+1,j}h_{n-1,j+1} + \cdots \\ &= h_{n-1,j}h_{n+1,j-1} + h_{n+1,j}h_{n-1,j+1} + \cdots, \\ \text{so } d(h_{2n,j} + h_{2n,j+1}) &= h_{n-1,j}h_{n+1,j-1} + h_{n+1,j}h_{n-1,j+1} \\ &\quad h_{n-1,j+1}h_{n+1,j} + h_{n+1,j+1}h_{n-1,j+2} + \cdots \\ &= h_{n-1,j}h_{n+1,j-1} + h_{n+1,j+1}h_{n-1,j+2} + \cdots, \end{aligned}$$

and so on.

Finally we need to consider the cases of (2.6) where  $m + i$  is not divisible by  $n$ . For this we can show that the expressions

$$\sum_{m < k < i} h_{k,j}h_{m+i-k,j+k}$$

are linearly independent. Suppose the term

$$\pm h_{k,x}h_{m+i-k,y}$$

appears the sums for some value of  $j$ . Then modulo  $n$  either  $j = x$  and  $y \equiv k + x$ , so  $x \equiv y - k$ , or  $j = y$  and  $x \equiv m + i + y - k$ . These conditions on  $x$  are mutually exclusive since  $m + i$  is not divisible by  $n$ . This means that each monomial of this form can appear in the sum for at most one value of  $j$ , so the sums for various  $j$  are linearly independent.  $\square$

Now  $\text{Ext}_{S(n,m+1)}^1$  is a subspace of  $H^1(L(n, m+1))$ . To finish the proof of the theorem we need to show that the elements  $h_{m+i,j}$  with  $i > \max(c-m, m+1)$  do not survive passage to  $\text{Ext}_{E^0S(n,m+1)}^1$  or from it to  $\text{Ext}_{S(n,m+1)}^1$ . We need to look at the first and second spectral sequences constructed for this purpose by May in [May66] and described (for  $m = 0$ ) in [Rav86, 6.3.4]. It follows from (2.2) that in the first May spectral sequence

$$d_r(h_{m+i,j}) = b_{m+i-n,j-1} \neq 0 \quad \text{for } i > n$$

for some  $r$ .

This eliminates all of the unwanted primitives except the ones with

$$\max(c-m, m+1) < i \leq n.$$

For this we can use (2.1), which implies that in the second May spectral sequence,

$$d_r(h_{m+i,j}) = \sum_{m < k < i} h_{k,j}h_{m+i-k,j+k}$$

where

$$\begin{aligned} r &= \min(d_{n,m+i} - d_{n,k} - d_{n,m+i-k} : m < k < i) \\ &= p(m+i-n) - (m+i) \\ &\quad \text{since } k \text{ and } m-i-k \text{ do not exceed } n \text{ and } m+i < 2n \\ &= (p-1)(m+i) - pn. \end{aligned}$$

Note that

$$n < c < m+i \leq m+n < 2n$$

so  $m + i$  is not divisible by  $n$ . Thus we can argue as in the last paragraph of the proof of Lemma 2.4 that the sums  $\sum_{m < k < i} h_{k,j} h_{m+i-k,j+k}$  are linearly independent. It follows that no linear combination of the unwanted  $h_{m+i,j}$  can survive to  $\text{Ext}_{S(n,m+1)}^1$ , so  $\text{Ext}_{\Sigma(n,m+1)}^1$  is as claimed.

## REFERENCES

- [May66] J. P. May. The cohomology of restricted Lie algebras and of Hopf algebras. *Journal of Algebra*, 3:123–146, 1966.
- [MM65] J. W. Milnor and J. C. Moore. On the structure of Hopf algebras. *Annals of Mathematics*, 81(2):211–264, 1965.
- [MR77] H. R. Miller and D. C. Ravenel. Morava stabilizer algebras and the localization of Novikov’s  $E_2$ -term. *Duke Mathematical Journal*, 44:433–447, 1977.
- [MRW77] H. R. Miller, D. C. Ravenel, and W. S. Wilson. Periodic phenomena in the Adams-Novikov spectral sequence. *Annals of Mathematics*, 106:469–516, 1977.
- [Rav86] D. C. Ravenel. *Complex Cobordism and Stable Homotopy Groups of Spheres*. Academic Press, New York, 1986. Errata available at <http://www.math.rochester.edu/u/drav/mu.html>.

UNIVERSITY OF ROCHESTER, ROCHESTER, NY 14627