

The Segal conjecture for topological Hochschild homology of complex cobordism

Sverre Lunøe-Nielsen and John Rognes

ABSTRACT

We study the C_p -equivariant Tate construction on the topological Hochschild homology $\mathrm{THH}(B)$ of a symmetric ring spectrum B by relating it to a topological version $R_+(B)$ of the Singer construction, extended by a natural circle action. This enables us to prove that the fixed and homotopy fixed point spectra of $\mathrm{THH}(B)$ are p -adically equivalent for $B = \mathrm{MU}$ and BP . This generalizes the classical C_p -equivariant Segal conjecture, which corresponds to the case $B = S$.

Contents

1. Introduction	591
2. Strategy of proof	592
3. Equivariant approximations	595
4. Tate representatives	601
5. Cyclotomic structure	608
6. The comparison map	610
7. The Segal conjecture	615
References	621

1. Introduction

Let p be a prime, let $\mathbb{T}$ be the circle group and let $C_p \subset \mathbb{T}$ be the cyclic subgroup of order p . Let B be any symmetric ring spectrum, in the sense of [20]. Its topological Hochschild homology $\mathrm{THH}(B)$, as constructed in [19], is then a genuinely $\mathbb{T}$ -equivariant spectrum in the sense of [21]. Let $E\mathbb{T}$ be a free, contractible $\mathbb{T}$ -CW complex and let $c: E\mathbb{T}_+ \rightarrow S^0$ be the usual collapse map. Forming the induced map $\mathrm{THH}(B) = F(S^0, \mathrm{THH}(B)) \rightarrow F(E\mathbb{T}_+, \mathrm{THH}(B))$ and passing to C_p -fixed points, we obtain the canonical map $\Gamma: \mathrm{THH}(B)^{C_p} \rightarrow \mathrm{THH}(B)^{hC_p}$ comparing fixed and homotopy fixed points.

When $B = S$ is the sphere spectrum, $\mathrm{THH}(S)$ and the $\mathbb{T}$ -equivariant sphere spectrum are C_p -equivariantly equivalent, so the Segal conjecture for C_p is the assertion that $\Gamma: \mathrm{THH}(S)^{C_p} \rightarrow \mathrm{THH}(S)^{hC_p}$ is a p -adic equivalence, which was proved in [2, 22]. Our main theorem generalizes this result to the case when $B = \mathrm{MU}$ is the complex cobordism spectrum or $B = \mathrm{BP}$ is the p -local Brown–Peterson spectrum.

THEOREM 1.1. *The maps*

$$\Gamma: \mathrm{THH}(\mathrm{MU})^{C_p} \longrightarrow \mathrm{THH}(\mathrm{MU})^{hC_p}$$

and

$$\Gamma: \mathrm{THH}(\mathrm{BP})^{C_p} \longrightarrow \mathrm{THH}(\mathrm{BP})^{hC_p}$$

are p -adic equivalences.

Received 27 October 2010; revised 21 April 2011.

2010 *Mathematics Subject Classification* 55P91 (primary), 55P43, 55S10 (secondary).

The corresponding assertion is not true as stated for general symmetric ring spectra B , but there are special cases for which it is approximately true. For example, when $B = H\mathbb{F}_p, H\mathbb{Z}$ (the Eilenberg–Mac Lane spectra) or ℓ (the Adams summand of connective topological K -theory ku), the map Γ induces an isomorphism of homotopy groups in sufficiently high degrees, with suitable coefficients, as proved in [19, 12, 4] respectively. In particular, in those cases the homotopy fixed point spectrum is not connective, even if the fixed point spectrum is. We believe that the explanation for this phenomenon is related to the conjectured ‘red-shift’ property of topological cyclic homology and algebraic K -theory [5]. In the examples mentioned above, the difference between the fixed and homotopy fixed points stems from the first chromatic truncation present in B (a Milnor Q^i that acts trivially on $H^*(B; \mathbb{F}_p)$, followed by a Q^{i+1} that acts non-trivially), whereas S , MU and BP are not truncated at any finite chromatic complexity. The p -adic equivalences discussed above are thus rather special properties of the symmetric ring spectra S , MU and BP . This is reflected in our proof of Theorem 1.1, which depends on calculational facts particular to these cases.

Let $C_{p^n} \subset \mathbb{T}$ be the cyclic subgroup of order p^n . The original Segal conjecture for C_{p^n} is known to follow from the one for C_p , without further explicit computations [27], much as the Segal conjecture for general p -groups follows from the elementary abelian case [15]. Similarly, combining [10, 1.8] with Theorem 1.1 implies the following corollary.

COROLLARY 1.2. *The maps*

$$\Gamma_n: \mathrm{THH}(\mathrm{MU})^{C_{p^n}} \rightarrow \mathrm{THH}(\mathrm{MU})^{hC_{p^n}}$$

and

$$\Gamma_n: \mathrm{THH}(\mathrm{BP})^{C_{p^n}} \rightarrow \mathrm{THH}(\mathrm{BP})^{hC_{p^n}}$$

are p -adic equivalences, for all $n \geq 1$.

1.1. Organization

In § 2, we outline how to deduce the main theorem from the existence of a suitable bicontinuous isomorphism Φ_B for $B = MU$ or BP . This isomorphism is constructed in § 7, but this requires computational control of a canonical map $\gamma: \mathrm{THH}(B) \rightarrow \mathrm{THH}(B)^{tC_p}$. It suffices to gain this control on the image of a map $\omega: \mathbb{T} \ltimes B \rightarrow \mathrm{THH}(B)$. For this we use a diagram (2.4), which is constructed in § 5. Thus, $\gamma \circ \omega$ factors as the composite of a map $\rho \wedge \epsilon_B: \mathbb{T} \ltimes B \rightarrow \mathbb{T}/C_p \ltimes R_+(B)$, which was analysed in our paper [23], and a map $\omega^t: \mathbb{T}/C_p \ltimes R_+(B) \rightarrow \mathrm{THH}(B)^{tC_p}$, which is constructed and calculated in §§ 3 and 4. The calculations that are special to $B = MU$ and BP are completed in § 6.

1.2. History

This work started out as a part of the first author’s PhD thesis at the University of Oslo, supervised by the second author. The original thesis covered the case of BP at $p = 2$, under the assumption that BP is an E_∞ ring spectrum. The current version does not make that assumption, and covers both the cases MU and BP , at all primes.

2. Strategy of proof

In this section, we outline how to deduce the Main Theorem 1.1 from the cohomological Theorem 2.1 or, equivalently, from the homological Theorem 2.2.

Let $\widetilde{E\mathbb{T}}$ be the mapping cone of the collapse map $c: E\mathbb{T}_+ \rightarrow S^0$. There is a homotopy cartesian square of $\mathbb{T}$ -equivariant spectra

$$\begin{array}{ccc} \mathrm{THH}(B) & \longrightarrow & \widetilde{E\mathbb{T}} \wedge \mathrm{THH}(B) \\ \downarrow & & \downarrow \\ F(E\mathbb{T}_+, \mathrm{THH}(B)) & \longrightarrow & \widetilde{E\mathbb{T}} \wedge F(E\mathbb{T}_+, \mathrm{THH}(B)) \end{array}$$

see [18]. Passing to C_p -fixed points, we get a homotopy cartesian square of $\mathbb{T}/C_p$ -equivariant spectra

$$\begin{array}{ccc} \mathrm{THH}(B)^{C_p} & \xrightarrow{R} & [\widetilde{E\mathbb{T}} \wedge \mathrm{THH}(B)]^{C_p} \\ \Gamma \downarrow & & \downarrow \hat{\Gamma} \\ \mathrm{THH}(B)^{hC_p} & \xrightarrow{R^h} & \mathrm{THH}(B)^{tC_p}. \end{array} \quad (2.1)$$

Here $\mathrm{THH}(B)^{tC_p}$ denotes the C_p -Tate construction on $\mathrm{THH}(B)$. In order to prove Theorem 1.1, we use the homotopy cartesian square (2.1) to translate the problem into a question about the map $\hat{\Gamma}: [\widetilde{E\mathbb{T}} \wedge \mathrm{THH}(B)]^{C_p} \rightarrow \mathrm{THH}(B)^{tC_p}$.

Suppose hereafter that B is connective, meaning that $\pi_*(B) = 0$ for $* < 0$. By the cyclotomic structure of $\mathrm{THH}(B)$ (see [19]) there is a natural equivalence $[\widetilde{E\mathbb{T}} \wedge \mathrm{THH}(B)]^{C_p} \xrightarrow{\simeq} \mathrm{THH}(B)$, and we are faced with the problem of showing that the composite stable map

$$\gamma: \mathrm{THH}(B) \longrightarrow \mathrm{THH}(B)^{tC_p}$$

is a p -adic equivalence for $B = \mathrm{MU}$ and BP . We do this using the methods of [16, 23], by realizing the Tate construction $\mathrm{THH}(B)^{tC_p}$ as the homotopy limit

$$\mathrm{THH}(B)^{tC_p} \simeq \operatorname{holim}_n \mathrm{THH}(B)^{tC_p}[n]$$

of a Tate tower $\{\mathrm{THH}(B)^{tC_p}[n]\}_n$ (see (4.5)) of bounded below spectra of finite type mod p , and comparing the Adams spectral sequence

$$E_2^{s,t} = \operatorname{Ext}_{\mathcal{A}}^{s,t}(H^*(\mathrm{THH}(B)), \mathbb{F}_p) \Longrightarrow \pi_{t-s} \mathrm{THH}(B)_p^\wedge \quad (2.2)$$

to the inverse limit of Adams spectral sequences

$$E_2^{s,t} = \operatorname{Ext}_{\mathcal{A}}^{s,t}(H_c^*(\mathrm{THH}(B)^{tC_p}), \mathbb{F}_p) \Longrightarrow \pi_{t-s}(\mathrm{THH}(B)^{tC_p})_p^\wedge \quad (2.3)$$

associated to that tower. We always use $\mathbb{F}_p$ -coefficients for homology and cohomology, $\mathcal{A}$ denotes the Steenrod algebra and

$$H_c^*(\mathrm{THH}(B)^{tC_p}) = \operatorname{colim}_n H^*(\mathrm{THH}(B)^{tC_p}[n])$$

is the continuous cohomology of the Tate tower. We note, as in [23, 2.3], that this colimit depends on the actual tower, not just on its homotopy limit. The key computational input is now that, in the cases $B = S$, MU and BP , the induced $\mathcal{A}$ -module homomorphism

$$\gamma^*: H_c^*(\mathrm{THH}(B)^{tC_p}) \longrightarrow H^*(\mathrm{THH}(B))$$

is an Ext-equivalence, in the sense that it induces an isomorphism between the E_2 -terms of the spectral sequences (2.2) and (2.3). Both spectral sequences are strongly convergent for connective B of finite type mod p , so this implies that γ , $\hat{\Gamma}$ and Γ are p -adic equivalences. More precisely, we can prove the following theorem. Recall from [2, 23] that the Singer construction on an $\mathcal{A}$ -module M is an $\mathcal{A}$ -module $R_+(M)$, which comes equipped with a natural Ext-equivalence $\epsilon: R_+(M) \rightarrow M$.

THEOREM 2.1. *When $B = S$, MU or BP, there is an $\mathcal{A}$ -module isomorphism*

$$\Phi_B^*: H_c^*(\mathrm{THH}(B)^{tC_p}) \xrightarrow{\cong} R_+(H^*(\mathrm{THH}(B)))$$

such that $\gamma^* = \epsilon \circ \Phi_B^*$.

As summarized above, this implies Theorem 1.1. When $B = S$, the 0-simplex inclusion $\eta_p: S^{\wedge p} \rightarrow \mathrm{sd}_p \mathrm{THH}(S) \cong \mathrm{THH}(S)$ (see Definition 3.6) is a C_p -equivariant equivalence, so the theorem is already covered by Lunøe–Nielsen and Rognes [23, 5.13], or the original proofs of the Segal conjecture. On the other hand, the cases $B = \mathrm{MU}$ and BP involve new ideas. For one thing, we will make use of the ring spectrum structure on $\mathrm{THH}(B)^{tC_p}$, which means that it is more convenient to work with the *continuous homology*

$$H_*^c(\mathrm{THH}(B)^{tC_p}) = \lim_n H_*(\mathrm{THH}(B)^{tC_p}[n])$$

of the Tate tower, and its induced algebra structure, than with the continuous cohomology. The continuous homology must be viewed as a topological graded $\mathbb{F}_p$ -vector space, with the linear topology generated by the neighbourhood basis of the origin given by the kernels

$$F_n H_*^c(\mathrm{THH}(B)^{tC_p}) = \ker(H_*^c(\mathrm{THH}(B)^{tC_p}) \rightarrow H_*(\mathrm{THH}(B)^{tC_p}[n])).$$

We may also view $H_*^c(\mathrm{THH}(B)^{tC_p})$ as the limit of its quotients

$$F^n H_*^c(\mathrm{THH}(B)^{tC_p}) = \mathrm{im}(H_*^c(\mathrm{THH}(B)^{tC_p}) \rightarrow H_*(\mathrm{THH}(B)^{tC_p}[n])).$$

Let $\mathcal{A}_*$ denote the dual of the Steenrod algebra. The $\mathcal{A}_*$ -comodule structure on $H_*(\mathrm{THH}(B)^{tC_p}[n])$ for each integer n makes the limit $H_*^c(\mathrm{THH}(B)^{tC_p})$ a *complete $\mathcal{A}_*$ -comodule* in the sense of [23, 2.7]. Similarly, for each $\mathcal{A}_*$ -comodule M_* that is bounded below and of finite type, the *homological Singer construction* $R_+(M_*)$ from [23, 3.7] is a complete $\mathcal{A}_*$ -comodule, equipped with a natural continuous homomorphism $\epsilon_*: M_* \rightarrow R_+(M_*)$ of complete $\mathcal{A}_*$ -comodules. The linear topologies on $H_*^c(\mathrm{THH}(B)^{tC_p})$ and $R_+(M_*)$ allow us to recover the continuous cohomology and the (cohomological) Singer construction, respectively, as their continuous $\mathbb{F}_p$ -linear duals

$$\begin{aligned} H_c^*(\mathrm{THH}(B)^{tC_p}) &\cong \mathrm{Hom}^c(H_*^c(\mathrm{THH}(B)^{tC_p}), \mathbb{F}_p) \\ R_+(M) &\cong \mathrm{Hom}^c(R_+(M_*), \mathbb{F}_p), \end{aligned}$$

where $M = H^*(\mathrm{THH}(B))$ and $M_* = H_*(\mathrm{THH}(B))$; see [23, 2.6, 2.9]. The corresponding assertions for the full linear duals would be false, as neither $H_c^*(\mathrm{THH}(B)^{tC_p})$ nor $R_+(M)$ will be of finite type. The $\mathcal{A}$ -module homomorphism γ^* is the continuous dual of the complete $\mathcal{A}_*$ -comodule homomorphism

$$\gamma_*: H_*(\mathrm{THH}(B)) \longrightarrow H_*^c(\mathrm{THH}(B)^{tC_p}),$$

where $H_*(\mathrm{THH}(B))$ has the discrete topology, and similarly ϵ is the continuous dual of ϵ_* . Theorem 2.1 now follows immediately from its homological analogue, stated below, by letting Φ_B^* be the continuous dual of Φ_B .

THEOREM 2.2. *When $B = S$, MU or BP, there is a continuous isomorphism of complete $\mathcal{A}_*$ -comodules*

$$\Phi_B: R_+(H^*(\mathrm{THH}(B))) \xrightarrow{\cong} H_*^c(\mathrm{THH}(B)^{tC_p}),$$

with continuous inverse, such that $\gamma_* = \Phi_B \circ \epsilon_*$.

This is then the main technical result that we need to prove. The argument relies on computations, and is not of a formal nature. To effect these computations, we establish in

Theorem 5.3 that, for connective symmetric ring spectra B , there is a natural commutative square

$$\begin{array}{ccc} \mathbb{T} \ltimes B & \xrightarrow{\omega} & \mathrm{THH}(B) \\ \rho \ltimes \epsilon_B \downarrow & & \downarrow \gamma \\ \mathbb{T}/C_p \ltimes R_+(B) & \xrightarrow{\omega^t} & \mathrm{THH}(B)^{tC_p} \end{array} \quad (2.4)$$

in the stable homotopy category, where γ is as above, $\rho: \mathbb{T} \rightarrow \mathbb{T}/C_p$ is the p th root isomorphism of groups, $R_+(B) = (B^{\wedge p})^{tC_p}$ is the *topological Singer construction* from [23, 5.8] realizing the homological Singer construction in continuous homology and $\epsilon_B: B \rightarrow R_+(B)$ is a natural map inducing the continuous homomorphism $\epsilon_*: H_*(B) \rightarrow R_+(H_*(B)) \cong H_*^c(R_+(B))$ of complete $\mathcal{A}_*$ -comodules. The map ω extends the inclusion of 0-simplices $\eta: B \rightarrow \mathrm{THH}(B)$ using the $\mathbb{T}$ -action on the target, and the construction of ω^t is similar, but more elaborate.

To compute the continuous homology of $\mathrm{THH}(B)^{tC_p}$, we use the homological Tate spectral sequence

$$\widehat{E}_{*,*}^2 = \widehat{H}^{-*}(C_p; H_*(\mathrm{THH}(B))) \implies H_*^c(\mathrm{THH}(B)^{tC_p})$$

recalled in Proposition 4.1. In the cases we are interested in, $H_*(\mathrm{THH}(B))$ is generated as an algebra by the image of ω_* , so we can use the commutative square (2.4), known formulas [23, § 3.2.1] for ϵ_* and new explicit formulas established in Theorem 4.11 for ω_*^t , in order to understand the completed $\mathcal{A}_*$ -comodule homomorphism γ_* . These calculations are carried out for $B = \mathrm{MU}$ and BP in Propositions 6.3 and 6.8, and in Theorems 6.4 and 6.9. The proof of Theorem 2.2 is completed in § 7, the main steps being Propositions 7.2 and 7.3.

3. Equivariant approximations

In this section, we specify our model for $\mathrm{THH}(B)$ in Definition 3.1, and obtain a natural map

$$\omega_p: \mathbb{T} \ltimes_{C_p} B^{\wedge p} \rightarrow \mathrm{sd}_p \mathrm{THH}(B)$$

to its p -fold edgewise subdivision in Definition 3.6. In the subsequent lemmas, we analyse its effect in homology, and in the Bökstedt spectral sequence $HH_*(H_*(B)) \implies H_*(\mathrm{THH}(B))$. We assume some familiarity with equivariant spectra [21], symmetric ring spectra [20], edgewise subdivision of cyclic objects [11, § 1] and the topological Hochschild homology spectrum [19, § 2.4].

Recall Connes' extension Λ of the category Δ . A cyclic space $X_\bullet$ is a contravariant functor from Λ to spaces, and its geometric realization $|X_\bullet|$ has a natural $\mathbb{T}$ -action. The $\mathbb{T}$ -action on a 0-simplex x in X_0 traces out the closed loop in $|X_\bullet|$ given by the 1-simplex $t_1 s_0(x)$ in X_1 .

Let B be a symmetric ring spectrum, with n th space B_n for each $n \geq 0$, and let I be Bökstedt's category [9] of finite sets $\mathbf{n} = \{1, 2, \dots, n\}$ for $n \geq 0$ and injective functions.

DEFINITION 3.1. For each $k \geq 0$ and each finite-dimensional $\mathbb{T}$ -representation V , let

$$\mathrm{thh}(B; S^V)_k = \mathrm{hocolim}_{(\mathbf{n}_0, \dots, \mathbf{n}_k) \in I^{k+1}} \mathrm{Map}(S^{\mathbf{n}_0} \wedge \dots \wedge S^{\mathbf{n}_k}, B_{\mathbf{n}_0} \wedge \dots \wedge B_{\mathbf{n}_k} \wedge S^V).$$

These spaces combine to a cyclic space $\mathrm{thh}(B; S^V)_\bullet$, with Hochschild-style structure maps d_i , s_j and t_k . Its geometric realization $\mathrm{thh}(B; S^V) = |\mathrm{thh}(B; S^V)_\bullet|$ has a natural $\mathbb{T}$ -action, given as the diagonal of the $\mathbb{T}$ -action coming from the cyclic structure and the $\mathbb{T}$ -action on S^V . These $\mathbb{T}$ -spaces assemble to a $\mathbb{T}$ -equivariant prespectrum $\mathrm{thh}(B)$, with V th space $\mathrm{thh}(B)(V) = \mathrm{thh}(B; S^V)$. We let

$$\mathrm{THH}(B) = L(\mathrm{thh}(B)^\tau)$$

be the genuine $\mathbb{T}$ -spectrum obtained by spectrification from a functorial good thickening of this prespectrum. (See [19, §A.1] for the notion of a *good* prespectrum, and a functorial construction of such a good thickening.)

DEFINITION 3.2. Let $B_{\text{pre}}^{\wedge 1}$ be the prespectrum with V th space

$$B_{\text{pre}}^{\wedge 1}(V) = \text{thh}(B; S^V)_0 = \text{hocolim}_{\mathbf{n} \in I} \text{Map}(S^n, B_n \wedge S^V),$$

and let $B^{\wedge 1} = L((B_{\text{pre}}^{\wedge 1})^\tau)$. The inclusion of 0-simplices defines a natural map

$$\eta: B^{\wedge 1} \rightarrow \text{THH}(B)$$

of spectra. Using the $\mathbb{T}$ -action on the target, we can uniquely extend η to a map

$$\omega: \mathbb{T} \ltimes B^{\wedge 1} \rightarrow \text{THH}(B)$$

of $\mathbb{T}$ -equivariant spectra, as in [21, II.4.1].

We suppose hereafter that B is connective. Without loss of generality, we may also assume that B is flat and convergent, as defined in [23, §5.2]. Then the natural maps

$$B_{\text{pre}}^{\wedge 1}(V) \xleftarrow{\simeq} B_{\text{pre}}^{\wedge 1}(V)^\tau \xrightarrow{\simeq} B^{\wedge 1}(V)$$

and

$$\text{thh}(B)(V) \xleftarrow{\simeq} \text{thh}(B)(V)^\tau \xrightarrow{\simeq} \text{THH}(B)(V)$$

are C -equivariant equivalences for each finite subgroup $C \subset \mathbb{T}$, by Hesselholt and Madsen [19, Proposition 2.4]. Furthermore, there is a natural chain of weak equivalences relating $B^{\wedge 1}$ and the Lewis–May spectrum associated to B ; see [23, 5.5]. Hence, we shall simply write B for $B^{\wedge 1}$, especially in the context of the maps $\eta: B \rightarrow \text{THH}(B)$ and $\omega: \mathbb{T} \ltimes B \rightarrow \text{THH}(B)$.

For any simplicial space $X_\bullet$, its p -fold edgewise subdivision $\text{sd}_p X_\bullet$ is obtained by precomposing the contravariant functor $X_\bullet$ with the functor

$$\text{sd}_p: \Delta \rightarrow \Delta$$

that takes the object $[k] = \{0 < 1 < \dots < k\}$ to its p -fold concatenation

$$\text{sd}_p([k]) = [k] \sqcup \dots \sqcup [k] = [p(k+1)-1],$$

for each $k \geq 0$, and likewise for morphisms: $\text{sd}_p(f) = f \sqcup \dots \sqcup f$. Hence, $(\text{sd}_p X)_k = X_{p(k+1)-1}$. There is a natural homeomorphism [11, 1.1]

$$D: |\text{sd}_p X_\bullet| \xrightarrow{\cong} |X_\bullet|,$$

induced by the diagonal embeddings $\Delta^k \rightarrow \Delta^{p(k+1)-1}$ given in barycentric coordinates as

$$u \mapsto (u/p, \dots, u/p),$$

for $u = (u_0, \dots, u_k) \in \Delta^k$, with $\sum_i u_i = 1$, $u_i \geq 0$. In more detail, this is the map

$$D: \coprod_{k \geq 0} (\text{sd}_p X)_k \times \Delta^k / \sim \longrightarrow \coprod_{\ell \geq 0} X_\ell \times \Delta^\ell / \sim$$

that takes the k th summand to the ℓ th one, with $\ell = p(k+1) - 1$, via the identity $(\text{sd}_p X)_k = X_\ell$ and the given embedding $\Delta^k \rightarrow \Delta^\ell$.

DEFINITION 3.3. There is a natural simplicial map $e_\bullet: \text{sd}_p X_\bullet \rightarrow X_\bullet$, induced by the natural transformation $\text{id} \rightarrow \text{sd}_p$ of functors $\Delta \rightarrow \Delta$, with components $[k] \rightarrow \text{sd}_p([k])$ given

by inclusion on the p th (= last) copy of $[k]$ for $k \geq 0$. This is the order-preserving function that omits the $(p-1)(k+1)$ first elements in the target, and so, in simplicial degree k ,

$$e_k = d_0^{(p-1)(k+1)} : (\mathrm{sd}_p X)_k = X_{p(k+1)-1} \rightarrow X_k \quad (3.1)$$

is equal to the $(p-1)(k+1)$ -fold iterate of the 0th face map. After geometric realization,

$$e = |e_\bullet| : |\mathrm{sd}_p X_\bullet| \rightarrow |X_\bullet|$$

is induced by the corresponding iterated face maps $\delta_0^{(p-1)(k+1)} : \Delta^k \rightarrow \Delta^{p(k+1)-1}$, given in barycentric coordinates as

$$u \mapsto (0, \dots, 0, u),$$

for $u \in \Delta^k$, as above.

LEMMA 3.4. *There is a natural homotopy $D \simeq e = |e_\bullet|$ of maps $|\mathrm{sd}_p X_\bullet| \rightarrow |X_\bullet|$.*

Proof. The homotopy is given by the convex linear combination

$$(t, u) \mapsto (1-t)(u/p, \dots, u/p) + t(0, \dots, 0, u),$$

for $t \in [0, 1]$, $u \in \Delta^k$, of the two maps $\Delta^k \rightarrow \Delta^{p(k+1)-1}$. $\square$

Recall the ‘ p -fold cover’ $\Lambda_p \rightarrow \Lambda$ [11, 1.5]. Roughly, Λ_p is like Connes’ category Λ , except that the cyclic morphism $\tau_k : [k] \rightarrow [k]$ satisfies $\tau_k^{p(k+1)} = \mathrm{id}$ instead of $\tau_k^{k+1} = \mathrm{id}$. A Λ_p -space is a contravariant functor from Λ_p to spaces. When $X_\bullet$ is a cyclic space, with cyclic operators $t_k : X_k \rightarrow X_k$ satisfying $t_k^{k+1} = \mathrm{id}$ for $k \geq 0$, the p -fold edgewise subdivision $\mathrm{sd}_p X_\bullet$ is a Λ_p -space, with operators $t_k : (\mathrm{sd}_p X)_k \rightarrow (\mathrm{sd}_p X)_k$ satisfying $t_k^{p(k+1)} = \mathrm{id}$, defined to be equal to the operators $t_{p(k+1)-1} : X_{p(k+1)-1} \rightarrow X_{p(k+1)-1}$.

LEMMA 3.5. *Let $X_\bullet$ be a cyclic space. There is a natural $\mathbb{T}$ -action on $|\mathrm{sd}_p X_\bullet|$ such that $D : |\mathrm{sd}_p X_\bullet| \rightarrow |X_\bullet|$ is a $\mathbb{T}$ -equivariant homeomorphism. The action of the subgroup $C_p \subset \mathbb{T}$ is simplicial, in the sense that it comes from a C_p -action on $\mathrm{sd}_p X_\bullet$, such that a generator of C_p acts by*

$$t_k^{k+1} : (\mathrm{sd}_p X)_k \rightarrow (\mathrm{sd}_p X)_k,$$

for $k \geq 0$. In particular, the inclusion of 0-simplices $(\mathrm{sd}_p X)_0 \subset |\mathrm{sd}_p X_\bullet|$ is C_p -equivariant.

The $\mathbb{T}$ -action on a 0-simplex y in $(\mathrm{sd}_p X)_0 = X_{p-1}$ traces out the closed loop in $|\mathrm{sd}_p X_\bullet|$ given by the chain of p 1-simplices

$$t_1 s_0(y), t_1^3 s_0(y), \dots, t_1^{2p-1} s_0(y)$$

in $(\mathrm{sd}_p X)_1 = X_{2p-1}$. These meet at the p 0-simplices

$$y, t_1^2(y), \dots, t_1^{2p-2}(y)$$

that constitute the C_p -orbit of y .

Proof. See [11, 1.6] and the surrounding discussion. $\square$

DEFINITION 3.6. Let $\mathrm{sd}_p \mathrm{thh}(B) = |\mathrm{sd}_p \mathrm{thh}(B)_\bullet|$ be the $\mathbb{T}$ -equivariant prespectrum with V th space $\mathrm{sd}_p \mathrm{thh}(B)(V) = |\mathrm{sd}_p \mathrm{thh}(B; S^V)_\bullet|$, and let $\mathrm{sd}_p \mathrm{THH}(B) = L(\mathrm{sd}_p \mathrm{thh}(B)^\tau)$ be the spectrification of its good thickening, as in Definition 3.1. In simplicial degree 0,

$$\mathrm{sd}_p \mathrm{thh}(B; S^V)_0 = \mathrm{thh}(B; S^V)_{p-1} = B_{\mathrm{pre}}^{\wedge p}(V)$$

in the notation of [23, 5.3]. Hence, the inclusion of 0-simplices defines a natural map

$$\eta_p: B^{\wedge p} \rightarrow \mathrm{sd}_p \mathrm{THH}(B)$$

of C_p -equivariant spectra, where $B^{\wedge p} = L((B_{\mathrm{pre}}^{\wedge p})^\tau)$. Using the full $\mathbb{T}$ -action on the target, we can uniquely extend η_p to a map

$$\omega_p: \mathbb{T} \ltimes_{C_p} B^{\wedge p} \rightarrow \mathrm{sd}_p \mathrm{THH}(B)$$

of $\mathbb{T}$ -equivariant spectra, again as in [21, II.4.1].

Combining these constructions, we have a $\mathbb{T}$ -equivariant homeomorphism

$$D: \mathrm{sd}_p \mathrm{THH}(B) \xrightarrow{\cong} \mathrm{THH}(B).$$

LEMMA 3.7. *The composite*

$$D \circ \eta_p: B^{\wedge p} \rightarrow \mathrm{THH}(B)$$

is homotopic to the composite

$$e \circ \eta_p = \eta \circ d_0^{p-1}: B^{\wedge p} \rightarrow \mathrm{THH}(B),$$

where $d_0^{p-1}: B^{\wedge p} \rightarrow B^{\wedge 1} \simeq B$ is the p -fold multiplication map.

Proof. The homotopy is that of Lemma 3.4. The identity $e \circ \eta_p = \eta \circ d_0^{p-1}$ amounts to the formula (3.1) for e_k , in the special case $k = 0$. $\square$

To analyse the extension ω_p of η_p , we pass to homology.

DEFINITION 3.8. The circle $\mathbb{T}$, with the subgroup action of C_p , is a free C_p -CW complex. Its mod p cellular complex $C_*(\mathbb{T}) = \mathbb{F}_p[C_p]\{e_0, e_1\}$ has boundary operator $d(e_1) = (T - 1)e_0$, where $T \in C_p$ is the generator mapping to $\exp(2\pi i/p) \in \mathbb{T}$, e_1 is the 1-cell covering the arc from 1 to T and e_0 is the 0-cell 1.

Let $C_*(B)$ be the cellular complex of a CW-approximation to B , and choose a quasi-isomorphism $H_*(B) \simeq C_*(B)$, taking each homology class to a representing cycle in its class.

LEMMA 3.9. *There are quasi-isomorphisms*

$$C_*(\mathbb{T} \ltimes_{C_p} B^{\wedge p}) \simeq C_*(\mathbb{T}) \otimes_{C_p} C_*(B)^{\otimes p} \simeq C_*(\mathbb{T}) \otimes_{C_p} H_*(B)^{\otimes p},$$

and natural isomorphisms

$$\begin{aligned} H_*(\mathbb{T} \ltimes_{C_p} B^{\wedge p}) &\cong H_*(C_*(\mathbb{T}) \otimes_{C_p} H_*(B)^{\otimes p}) \\ &\cong \mathrm{cok}(T - 1)\{e_0\} \oplus \ker(T - 1)\{e_1\}, \end{aligned}$$

where

$$T - 1: H_*(B)^{\otimes p} \rightarrow H_*(B)^{\otimes p}$$

is the difference between the cyclic permutation, induced by the action of $T \in C_p$ on $B^{\wedge p}$, and the identity.

Proof. The first quasi-isomorphism follows from [21, VIII.1.6] and [21, VIII.2.9], combined with cellular approximation. We use [23, 5.5] to compare $B^{\wedge p}$ with the p th external power $B^{(p)}$. The second quasi-isomorphism is then clear, as $C_*(\mathbb{T})$ is C_p -free. This implies the first isomorphism. The second isomorphism is simply the computation of the homology of the complex

$$d: H_*(B)^{\otimes p}\{e_1\} \rightarrow H_*(B)^{\otimes p}\{e_0\},$$

where d maps $e_1 \otimes (\alpha_1 \otimes \dots \otimes \alpha_p)$ to $e_0 \otimes (T-1)(\alpha_1 \otimes \dots \otimes \alpha_p)$. $\square$

DEFINITION 3.10. For $\alpha \in H_q(B)$ let $e_1 \otimes \alpha^{\otimes p} \in H_{1+pq}(\mathbb{T} \ltimes_{C_p} B^{\wedge p})$ denote the homology class of $e_1 \otimes \alpha^{\otimes p}$ in $C_*(\mathbb{T}) \otimes_{C_p} H_*(B)^{\otimes p}$, and let

$$\alpha^{p-1} \wedge \alpha = (D \circ \omega_p)_*(e_1 \otimes \alpha^{\otimes p}) \in H_{1+pq}(\mathrm{THH}(B))$$

denote its image under $D \circ \omega_p: \mathbb{T} \ltimes_{C_p} B^{\wedge p} \rightarrow \mathrm{THH}(B)$.

The following spectral sequence was used by Bökstedt [9] to compute $\mathrm{THH}(\mathbb{Z}/p)$ and $\mathrm{THH}(\mathbb{Z})$. See [25, 3.1], [3, § 4] for more background.

LEMMA 3.11. *In the Bökstedt spectral sequence*

$$E_{*,*}^2 = HH_*(H_*(B)) \implies H_*(\mathrm{THH}(B)),$$

the class $\alpha^{p-1} \wedge \alpha$ is represented modulo filtration 0 by the homology class of the Hochschild 1-cycle

$$\alpha^{p-1} \otimes \alpha \in H_*(B) \otimes H_*(B).$$

Proof. By Lemma 3.4 we may compute $\alpha^{p-1} \wedge \alpha$ as the image of $e_1 \otimes \alpha^{\otimes p}$ under the cellular map $e \circ \omega_p$. By the second part of Lemma 3.5, and the formula (3.1) for $e = |e_\bullet|$ in simplicial degree $k = 1$, the cycle $e_1 \otimes \alpha^{\otimes p}$ maps under $e \circ \omega_p$ to the cycle

$$d_0^{2p-2} t_1 s_0(\alpha^{\otimes p}) = d_0^{2p-2}(1 \otimes \alpha \otimes \dots \otimes 1 \otimes \alpha) = \alpha^{p-1} \otimes \alpha$$

in the Bökstedt spectral sequence E^1 -term. $\square$

We can be more specific about $\alpha^{p-1} \wedge \alpha$ under additional commutativity hypotheses. The homotopy cofibre sequence

$$B \xrightarrow{i} \mathbb{T} \ltimes B \xrightarrow{j} S^1 \wedge B$$

is stably split by the collapse map $c: \mathbb{T} \ltimes B \rightarrow B$, and its homotopy fibre map, a section $s: S^1 \wedge B \rightarrow \mathbb{T} \ltimes B$. We let $\sigma: S^1 \wedge B \rightarrow \mathrm{THH}(B)$ be the composite map

$$S^1 \wedge B \xrightarrow{s} \mathbb{T} \ltimes B \xrightarrow{\omega} \mathrm{THH}(B).$$

For $\alpha \in H_q(B)$, the homology class $\sigma\alpha = \sigma_*(\alpha) \in H_{1+q}(\mathrm{THH}(B))$ is represented by the Hochschild 1-cycle $1 \otimes \alpha \in H_*(B) \otimes H_*(B)$ in the Bökstedt spectral sequence above; see [25, 3.2].

When B is an E_2 symmetric ring spectrum, it follows from [13] that $\mathrm{THH}(B)$ is an $E_1 = A_\infty$ ring spectrum, and hence equivalent to an (associative) S -algebra. Furthermore, the Bökstedt spectral sequence is an algebra spectral sequence with product induced by the shuffle product on the Hochschild complex. See, for example, [3, 4.2].

LEMMA 3.12. *Suppose that B is an E_2 symmetric ring spectrum. Then, for $\alpha \in H_q(B)$, we have*

$$\alpha^{p-1} \wedge \alpha \equiv \alpha^{p-1} \cdot \sigma \alpha$$

in $H_(\mathrm{THH}(B))$, modulo the image of $\eta_*: H_*(B) \rightarrow H_*(\mathrm{THH}(B))$.*

Proof. The shuffle product of the 0-cycle α^{p-1} and the 1-cycle $1 \otimes \alpha$ is the 1-cycle $\alpha^{p-1} \otimes \alpha$. Hence, the product $\alpha^{p-1} \cdot \sigma \alpha$ and the class $\alpha^{p-1} \wedge \alpha$ have the same representative in Hochschild filtration 1, and must be equal modulo the image of Hochschild filtration 0. $\square$

The following corollary applies to $B = S$, MU and BP. (According to Basterra and Mandell [7, 8], BP is an E_4 ring spectrum.)

COROLLARY 3.13. *Suppose that B is an E_2 symmetric ring spectrum with $H_*(B)$ concentrated in even degrees. Then, for $\alpha \in H_q(B)$, we have*

$$\alpha^{p-1} \wedge \alpha = \alpha^{p-1} \cdot \sigma \alpha.$$

Proof. There is only something to prove when q is even, in which case $H_{1+pq}(B) = 0$ by hypothesis, so the indeterminacy is 0. $\square$

When B is a commutative symmetric ring spectrum, it follows similarly that $\mathrm{THH}(B)$ is an E_∞ ring spectrum, hence equivalent to a commutative S -algebra. The multiplication maps $\mathrm{THH}(B)_k \simeq B^{\wedge(k+1)} \rightarrow B$ then combine to an augmentation map $\epsilon: \mathrm{THH}(B) \rightarrow B$ (not to be confused with the stable map ϵ_B), such that $\epsilon\eta = \mathrm{id}_B$ and $\epsilon\sigma \simeq *$. See, for example, [3, §3]. We can thus use ϵ to determine the error term in Lemma 3.12.

The commutative product on B makes the associated Lewis–May spectrum an E_∞ ring spectrum, with structure maps

$$\xi_p: E\Sigma_p \ltimes_{\Sigma_p} B^{(p)} \longrightarrow B$$

among others. Here $E\Sigma_p$ is a free, contractible Σ_p -space. Up to homotopy, there is a unique map $\chi: \mathbb{T} \rightarrow E\Sigma_p$ that is equivariant with respect to the inclusion $C_p \subset \Sigma_p$. Furthermore, the composite map

$$\mathbb{T} \ltimes_{C_p} B^{(p)} \xrightarrow{\chi \ltimes \mathrm{id}} E\Sigma_p \ltimes_{\Sigma_p} B^{(p)} \xrightarrow{\xi_p} B$$

exhibits the commuting homotopy $\mu^p \simeq \mu^p T$ that is preferred by the E_∞ structure, where $\mu^p: B^{(p)} \rightarrow B$ is the multiplication map and $T: B^{(p)} \rightarrow B^{(p)}$ is the cyclic permutation. Using [23, 5.5], we can identify the source here with the source $\mathbb{T} \ltimes_{C_p} B^{\wedge p}$ of ω_p .

LEMMA 3.14. *Let B be a commutative symmetric ring spectrum. The composite map*

$$\mathbb{T} \ltimes_{C_p} B^{\wedge p} \xrightarrow{\omega_p} \mathrm{sd}_p \mathrm{THH}(B) \xrightarrow{D} \mathrm{THH}(B) \xrightarrow{\epsilon} B$$

is homotopic to the composite map

$$\mathbb{T} \ltimes_{C_p} B^{\wedge p} \simeq \mathbb{T} \ltimes_{C_p} B^{(p)} \xrightarrow{\chi \ltimes \mathrm{id}} E\Sigma_p \ltimes_{\Sigma_p} B^{(p)} \xrightarrow{\xi_p} B.$$

Proof. Using Lemma 3.4, we may replace D by the simplicial map $e = |e_\bullet|$. When restricted to $1 \in C_p \subset \mathbb{T}$, both maps then agree with the p -fold multiplication map $d_0^{p-1}: B^{\wedge p} \rightarrow B$, taking $\alpha_1 \otimes \dots \otimes \alpha_p$ to $\alpha_1 \dots \alpha_p$ (in homology). Similarly, when restricted to $T \in C_p \subset \mathbb{T}$,

both maps agree with the cyclically permuted multiplication map $d_0^{p-1}t_{p-1}: B^{\wedge p} \rightarrow B$, taking $\alpha_1 \otimes \dots \otimes \alpha_p$ to $\alpha_p \alpha_1 \dots \alpha_{p-1}$. When restricted to the 1-cell $e_1 \subset \mathbb{T}$, connecting 1 to T , both maps exhibit the commuting homotopy between these two maps that is preferred by the given E_∞ structure. (This is how the E_∞ structure enters into the definition of ϵ .) In particular, the homotopies agree, up to homotopy relative to the end points. $\square$

The following corollary applies to $B = H\mathbb{F}_p, H\mathbb{Z}, ku$ and ℓ .

COROLLARY 3.15. *Let B be a commutative symmetric ring spectrum. For $\alpha \in H_q(B)$ we have*

$$\alpha^{p-1} \wedge \alpha = \alpha^{p-1} \cdot \sigma\alpha + \eta_*(Q_1(\alpha))$$

in $H_*(\mathrm{THH}(B))$, where $Q_1(\alpha) = \xi_{p*}(e_1 \otimes \alpha^{\otimes p})$ is the image of α under the Dyer–Lashof operation $Q_1: H_q(B) \rightarrow H_{1+pq}(B)$.

Proof. We know that $\alpha^{p-1} \wedge \alpha = (D\omega_p)_*(e_1 \otimes \alpha^{\otimes p})$ equals $\alpha^{p-1} \cdot \sigma\alpha + \eta_*(x)$ for some $x \in H_*(B)$. Applying ϵ_* , we get that $\epsilon_*(\alpha^{p-1} \wedge \alpha) = (\epsilon D\omega_p)_*(e_1 \otimes \alpha^{\otimes p}) = \xi_{p*}(e_1 \otimes \alpha^{\otimes p}) = Q_1(\alpha)$ equals $\epsilon_*(\alpha^{p-1} \cdot \sigma\alpha + \eta_*(x)) = \alpha^{p-1} \cdot 0 + x = x$. $\square$

4. Tate representatives

To prove Theorem 2.2, we use the homological Tate spectral sequence in Proposition 4.1 to study the continuous homology of Tate constructions. We only need to consider the C_p -equivariant case. Using Proposition 4.9, we deduce formulas in Theorem 4.11 for the effect of a natural map

$$\omega^t: \mathbb{T}/C_p \ltimes (B^{\wedge p})^{tC_p} \rightarrow \mathrm{THH}(B)^{tC_p},$$

as seen by the eyes of the Tate spectral sequence. These formulas will be essential for the calculations in § 6.

PROPOSITION 4.1. *Let X be a C_p -equivariant spectrum. Assume that X is bounded below with $H_*(X)$ of finite type. Then the homological Tate spectral sequence*

$$\widehat{E}_{s,t}^2(X) = \widehat{H}^{-s}(C_p; H_t(X)) \implies H_{s+t}^c(X^{tC_p})$$

converges strongly to the continuous homology of X^{tC_p} as a complete $\mathcal{A}_*$ -comodule.

If, furthermore, X is a C_p -equivariant ring spectrum, then this is an algebra spectral sequence, whose product at the $\widehat{E}^2$ -term is given by the cup product in Tate cohomology and the Pontryagin product on $H_*(X)$.

Proof. See Propositions 4.15 and 4.17 in [23]. $\square$

REMARK 4.2. The strong convergence of the spectral sequence refers to the complete Hausdorff filtration of the abutment $H_*^c(X^{tC_p})$ by the kernels of the natural homomorphisms induced by the maps $X^{tC_p} \rightarrow X^{tC_p}[n]$; see (4.5). We shall refer to this increasing filtration on $H_*^c(X^{tC_p})$ as the *Tate filtration*.

When B is a connective symmetric ring spectrum, with $H_*(B)$ of finite type, then both $X = B^{\wedge p}$ and $X = \mathrm{THH}(B)$ are C_p -spectra that are bounded below with $H_*(X)$ of finite type. This is clear from the Künneth formula and the Bökstedt spectral sequence in Lemma 3.11.

REMARK 4.3. In the case when $X = S^{\wedge p} \cong \mathrm{THH}(S)$ is the C_p -equivariant sphere spectrum, $H_*(X) = \mathbb{F}_p$ is concentrated in degree 0, so

$$\widehat{E}_{*,*}^2(S^{\wedge p}) = \widehat{H}^{-*}(C_p; \mathbb{F}_p)$$

is concentrated on the horizontal axis. Here $\widehat{H}^{-*}(C_p; \mathbb{F}_p) = P(u, u^{-1})$ for $p = 2$ and $\widehat{H}^{-*}(C_p; \mathbb{F}_p) = E(u) \otimes P(t, t^{-1})$ for p odd, with $u \in \widehat{H}^1$ and $t \in \widehat{H}^2$. There cannot be any differentials in this spectral sequence for bidegree reasons, so each class $u^i t^r$ is an infinite cycle. (This formula applies when p is odd; the reader should always replace $u^i t^r$ by u^{i+2r} when $p = 2$.)

By naturality with respect to the unit maps $S^{\wedge p} \rightarrow B^{\wedge p}$ and $\mathrm{THH}(S) \rightarrow \mathrm{THH}(B)$, it follows that the classes $u^i t^r$ are infinite cycles, also in the homological Tate spectral sequences for $X = B^{\wedge p}$ and $X = \mathrm{THH}(B)$. Hence, all of these spectral sequences exhibit a horizontal periodicity, with each $\widehat{E}^r$ -term being determined by the part on the vertical axis via an isomorphism

$$\widehat{E}_{*,*}^r(X) \cong \widehat{H}^{-*}(C_p; \mathbb{F}_p) \otimes \widehat{E}_{0,*}^r(X).$$

REMARK 4.4. In the case $X = B^{\wedge p}$ the C_p -action on $H_*(X) = H_*(B)^{\otimes p}$ is given by cyclic permutation of the tensor factors, so

$$\begin{aligned} \widehat{E}_{*,*}^2(B^{\wedge p}) &= \widehat{H}^{-*}(C_p; H_*(B)^{\otimes p}) \cong \widehat{H}^{-*}(C_p; \mathbb{F}_p) \otimes \mathbb{F}_p\{\alpha^{\otimes p}\} \\ &\implies H_*^c((B^{\wedge p})^{tC_p}), \end{aligned}$$

where α ranges over an $\mathbb{F}_p$ -basis of $H_*(B)$. Also in this case the spectral sequence collapses at the $\widehat{E}^2$ -term, and converges to

$$H_*^c((B^{\wedge p})^{tC_p}) = H_*^c(R_+(B)) \cong R_+(H_*(B));$$

see [23, 5.14]. Here $R_+(B) = (B^{\wedge p})^{tC_p}$ is the topological Singer construction on B , and $R_+(H_*(B))$ is the homological Singer construction on $H_*(B)$, discussed in [23, Definitions 5.8 and 3.7] respectively.

The right-hand isomorphism is given in Theorem 5.9 of that paper. Implicit in this isomorphism is the fact that, for each $u^i t^r \in \widehat{H}^{-*}(C_p; \mathbb{F}_p)$ and $\alpha \in H_*(B)$, there is a preferred class in $H_*^c((B^{\wedge p})^{tC_p})$ that is represented in the Tate spectral sequence by $u^i t^r \otimes \alpha^{\otimes p}$ in $\widehat{E}_{*,*}^2(B^{\wedge p})$. It is obtained by representing α by a map $f: S^q \rightarrow H \wedge B$, and taking $u^i t^r \otimes \alpha^{\otimes p}$ to be in the image of an induced homomorphism

$$H_*^c((f^p)^{tC_p}): H_*^c((S^{pq})^{tC_p}) \rightarrow H_*^c((B^{\wedge p})^{tC_p});$$

see the proof of [23, 5.14] for details. Hence, $u^i t^r \otimes \alpha^{\otimes p}$ is a well-defined class in $H_*^c((B^{\wedge p})^{tC_p})$, not just defined modulo the Tate filtration.

REMARK 4.5. In the case $X = \mathrm{THH}(B)$, the C_p -action on $H_*(X) = H_*(\mathrm{THH}(B))$ is obtained by restriction from a $\mathbb{T}$ -action, and hence is algebraically trivial. Thus,

$$\begin{aligned} \widehat{E}_{*,*}^2(\mathrm{THH}(B)) &= \widehat{H}^{-*}(C_p; H_*(\mathrm{THH}(B))) \cong \widehat{H}^{-*}(C_p; \mathbb{F}_p) \otimes H_*(\mathrm{THH}(B)) \\ &\implies H_*^c(\mathrm{THH}(B)^{tC_p}). \end{aligned}$$

In this case, the spectral sequence does not generally collapse at the $\widehat{E}^2$ -term. For example, the d^2 -differential satisfies

$$d^2(u^i t^r \otimes \alpha) = u^i t^{r+1} \otimes \sigma \alpha,$$

for p odd, and similarly for $p = 2$; see [29, 3.3]. When B is an E_∞ ring spectrum, so that $\mathrm{THH}(B)$ is equivalent to a $\mathbb{T}$ -equivariant commutative S -algebra, there are many cases where this spectral sequence collapses at the $\widehat{E}^3$ -term; see [14, 1.2]. For example, this is the case for $B = \mathrm{MU}$, as we shall prove directly in Proposition 6.3.

By naturality with respect to the C_p -equivariant map $D \circ \eta_p: B^{\wedge p} \rightarrow \mathrm{THH}(B)$, we get a map of homological Tate spectral sequences

$$\widehat{E}_{*,*}^2(B^{\wedge p}) \longrightarrow \widehat{E}_{*,*}^2(\mathrm{THH}(B))$$

taking $u^i t^r \otimes \alpha^{\otimes p}$ to $u^i t^r \otimes \alpha^p$ (recall Lemma 3.7), which converges to a complete $\mathcal{A}_*$ -comodule homomorphism

$$\eta_*^t: H_*^c((B^{\wedge p})^{tC_p}) \longrightarrow H_*^c(\mathrm{THH}(B)^{tC_p}).$$

This is useful for determining differentials and $\mathcal{A}_*$ -comodule structure in the target, but there is an even more useful extension of this homomorphism,

$$\omega_*^t: H_*(\mathbb{T}/C_p) \otimes H_*^c((B^{\wedge p})^{tC_p}) \longrightarrow H_*^c(\mathrm{THH}(B)^{tC_p}),$$

which is constructed by taking the full $\mathbb{T}$ -action on $\mathrm{THH}(B)$ into account. We now explain this construction.

Consider the diagram

$$\begin{array}{ccc} & \eta_p & \\ \curvearrowright & & \searrow \\ B^{\wedge p} & \xrightarrow{i_p} \mathbb{T} \ltimes_{C_p} B^{\wedge p} & \xrightarrow{\omega_p} \mathrm{sd}_p \mathrm{THH}(B) \\ & & \downarrow \cong D \\ & & \mathrm{THH}(B) \end{array} \quad (4.1)$$

of C_p -equivariant spectra, where i_p is induced by the inclusion $C_p \subset \mathbb{T}$. Note that ω_p and D are $\mathbb{T}$ -equivariant maps. For any closed subgroup $G \subseteq \mathbb{T}$ containing C_p , and any G -spectrum X , the C_p -Tate construction

$$X^{tC_p} = [\widetilde{E\mathbb{T}} \wedge F(E\mathbb{T}_+, X)]^{C_p}$$

is naturally a G/C_p -spectrum. Applying the C_p -Tate construction to (4.1), we can form a commutative diagram

$$\begin{array}{ccccc} & & \mathbb{T}/C_p \ltimes (B^{\wedge p})^{tC_p} & \xrightarrow{\omega^t} & \\ & \curvearrowright & \downarrow \simeq \kappa & & \\ (B^{\wedge p})^{tC_p} & \xrightarrow{i_p^{tC_p}} & (\mathbb{T} \ltimes_{C_p} B^{\wedge p})^{tC_p} & \xrightarrow{\omega_p^{tC_p}} & \mathrm{sd}_p \mathrm{THH}(B)^{tC_p} \\ & \searrow & & \downarrow \cong D^{tC_p} & \\ & & & & \mathrm{THH}(B)^{tC_p} \end{array} \quad (4.2)$$

η^t (curved arrow from $(B^{\wedge p})^{tC_p}$ to $\mathrm{THH}(B)^{tC_p}$)

DEFINITION 4.6. Let

$$\eta^t: (B^{\wedge p})^{tC_p} \rightarrow \mathrm{THH}(B)^{tC_p}$$

be the map $\eta^t = (D \circ \eta_p)^{tC_p}$, let

$$\kappa: \mathbb{T}/C_p \ltimes (B^{\wedge p})^{tC_p} \rightarrow (\mathbb{T} \ltimes_{C_p} B^{\wedge p})^{tC_p}$$

be the unique $\mathbb{T}/C_p$ -equivariant map that extends $i_p^{tC_p}$ and let

$$\omega^t: \mathbb{T}/C_p \ltimes (B^{\wedge p})^{tC_p} \rightarrow \mathrm{THH}(B)^{tC_p}$$

be the $\mathbb{T}/C_p$ -equivariant composite $\omega^t = (D \circ \omega_p)^{tC_p} \circ \kappa$.

LEMMA 4.7. *The map*

$$\kappa: \mathbb{T}/C_p \ltimes (B^{\wedge p})^{tC_p} \xrightarrow{\simeq} (\mathbb{T} \ltimes_{C_p} B^{\wedge p})^{tC_p}$$

is an equivalence.

Proof. The C_p -equivariant cofibre sequence

$$C_{p+} \xrightarrow{i_p} \mathbb{T}_+ \xrightarrow{j_p} S^1 \wedge C_{p+} \quad (4.3)$$

induces a map of cofibre sequences

$$\begin{array}{ccccc} (B^{\wedge p})^{tC_p} & \xrightarrow{i'} & \mathbb{T}/C_p \ltimes (B^{\wedge p})^{tC_p} & \longrightarrow & S^1 \wedge (B^{\wedge p})^{tC_p} \\ \downarrow = & & \downarrow \kappa & & \downarrow \simeq \\ (B^{\wedge p})^{tC_p} & \xrightarrow{i_p^{tC_p}} & (\mathbb{T} \ltimes_{C_p} B^{\wedge p})^{tC_p} & \longrightarrow & (S^1 \wedge B^{\wedge p})^{tC_p}, \end{array}$$

which implies the result. $\square$

REMARK 4.8. In the intermediate case of the homological Tate spectral sequence for $X = \mathbb{T} \ltimes_{C_p} B^{\wedge p}$, the homology $H_*(X) = H_*(\mathbb{T} \ltimes_{C_p} B^{\wedge p})$ was given in Lemma 3.9. The C_p -action extends to a $\mathbb{T}$ -action, and hence is algebraically trivial. Thus, in this case

$$\begin{aligned} \widehat{E}_{*,*}^2(\mathbb{T} \ltimes_{C_p} B^{\wedge p}) &= \widehat{H}^{-*}(C_p; H_*(\mathbb{T} \ltimes_{C_p} B^{\wedge p})) \cong \widehat{H}^{-*}(C_p; \mathbb{F}_p) \otimes H_*(\mathbb{T} \ltimes_{C_p} B^{\wedge p}) \\ &\implies H_*^c((\mathbb{T} \ltimes_{C_p} B^{\wedge p})^{tC_p}). \end{aligned}$$

It is not difficult to determine the d^2 -differentials, and to prove that this spectral sequence collapses at the $\widehat{E}^3$ -term, but we shall not need this result.

Recall the C_p -CW structure on $\mathbb{T}$ from Definition 3.8, which descends to a CW structure on the orbit space (= quotient group) $\mathbb{T}/C_p$, with mod p cellular complex $C_*(\mathbb{T}/C_p) = \mathbb{F}_p\{e_0, e_1\}$ having boundary operator $d(e_1) = 0$. Hence, $H_*(\mathbb{T}/C_p) = \mathbb{F}_p\{e_0, e_1\}$, with $\deg(e_j) = j$ for $j \in \{0, 1\}$.

PROPOSITION 4.9. *The map κ induces the complete $\mathcal{A}_*$ -comodule isomorphism*

$$\kappa_*: H_*(\mathbb{T}/C_p) \otimes H_*^c((B^{\wedge p})^{tC_p}) \xrightarrow{\cong} H_*^c((\mathbb{T} \ltimes_{C_p} B^{\wedge p})^{tC_p})$$

given, modulo Tate filtration in the target, by

$$e_j \otimes u^i t^r \otimes \alpha^{\otimes p} \longmapsto u^i t^r \otimes e_j \otimes \alpha^{\otimes p},$$

for $(i, j) = (0, 0), (1, 0)$ or $(0, 1)$, while

$$(e_1 \otimes u t^r - e_0 \otimes t^r) \otimes \alpha^{\otimes p} \longmapsto u t^r \otimes e_1 \otimes \alpha^{\otimes p}$$

in the case $(i, j) = (1, 1)$.

REMARK 4.10. Note that acting by the fundamental class of $\mathbb{T}/C_p$ on classes represented in odd filtration has the effect of increasing the Tate filtration by 1, as the image of $e_1 \otimes ut^r \otimes \alpha^{\otimes p}$ is represented by $t^r \otimes e_0 \otimes \alpha^{\otimes p}$, modulo a term of lower Tate filtration.

Proof. For definiteness, we assume that our models for $E\mathbb{T}$ and $\widetilde{E\mathbb{T}}$ are the unit sphere $S(\infty\mathbb{C})$ and the one-point compactification $S^{\infty\mathbb{C}}$ of $\infty\mathbb{C} = \mathbb{C}^\infty$ with the diagonal $\mathbb{T}$ -action respectively. We fix a based $\mathbb{T}$ -CW structure on $\widetilde{E\mathbb{T}}$ with $S^{n\mathbb{C}}$ as $\mathbb{T}$ -equivariant $(2n-1)$ - and $2n$ -skeleton, so that

$$S^{n\mathbb{C}}/S^{(n-1)\mathbb{C}} \cong \Sigma^{2n-1}\mathbb{T}_+.$$

Restricting the action to $C_p \subset \mathbb{T}$, there is a based C_p -CW structure on $\widetilde{E\mathbb{T}}$ with $S^{n\mathbb{C}}$ as C_p -equivariant $2n$ -skeleton, so that the subquotient C_p -CW structure on $S^{n\mathbb{C}}/S^{(n-1)\mathbb{C}}$ is the $(2n-1)$ th suspension of the C_p -CW structure on $\mathbb{T}$ from Definition 3.8, based at a disjoint base point. For $n \geq 0$ let $\widetilde{E}_n$ be the n -skeleton of this based C_p -CW structure on $\widetilde{E\mathbb{T}}$, viewed as a C_p -CW spectrum, and recall that the Greenlees filtration [17]

$$\cdots \rightarrow \widetilde{E}_{n-1} \rightarrow \widetilde{E}_n \rightarrow \cdots \rightarrow \widetilde{E\mathbb{T}}$$

of C_p -spectra extends this notation to all integers n , so that $\widetilde{E}_n/\widetilde{E}_{n-1} \cong \Sigma^n C_{p+}$ and $\widetilde{E}_{2n} = S^{n\mathbb{C}}$ for all $n \in \mathbb{Z}$. The cofibre sequence

$$\widetilde{E}_{2n-1}/\widetilde{E}_{2n-2} \rightarrow \widetilde{E}_{2n}/\widetilde{E}_{2n-2} \rightarrow \widetilde{E}_{2n}/\widetilde{E}_{2n-1} \quad (4.4)$$

equals the $(2n-1)$ th suspension of (4.3). If we only consider the even-indexed spectra, then we get a coarser filtration

$$\cdots \rightarrow \widetilde{E}_{2n-2} \rightarrow \widetilde{E}_{2n} \rightarrow \cdots \rightarrow \widetilde{E\mathbb{T}}$$

of $\mathbb{T}$ -spectra, so that $\widetilde{E}_{2n}/\widetilde{E}_{2n-2} \cong \Sigma^{2n-1}\mathbb{T}_+$ for all n .

The homological Tate spectral sequence in Proposition 4.1 is obtained by expressing the C_p -Tate construction as a homotopy limit

$$X^{tC_p} \simeq \operatorname{holim}_{n \rightarrow -\infty} X^{tC_p}[n]$$

of a tower of spectra, where

$$X^{tC_p}[n] = [\widetilde{E\mathbb{T}}/\widetilde{E}_{n-1} \wedge F(E\mathbb{T}_+, X)]^{C_p}. \quad (4.5)$$

There are homotopy (co-)fibre sequences

$$[\widetilde{E}_n/\widetilde{E}_{n-1} \wedge F(E\mathbb{T}_+, X)]^{C_p} \rightarrow X^{tC_p}[n] \rightarrow X^{tC_p}[n+1],$$

and equivalences

$$\begin{aligned} [\widetilde{E}_n/\widetilde{E}_{n-1} \wedge F(E\mathbb{T}_+, X)]^{C_p} &\simeq [\widetilde{E}_n/\widetilde{E}_{n-1} \wedge X]^{C_p} \\ &\simeq (\widetilde{E}_n/\widetilde{E}_{n-1} \wedge X)/C_p \cong \Sigma^n X, \end{aligned}$$

for all n , as $\widetilde{E}_n/\widetilde{E}_{n-1}$ is C_p -free. This uses the Adams transfer equivalence $E/C_p \simeq E^{C_p}$ for C_p -free spectra E ; see, for example, [21, § II.2]. The spectral sequence in question is associated to the exact couple with $A_{n,*} = H_{n+*}(X^{tC_p}[n])$ and

$$\widehat{E}_{n,*}^1 = H_{n+*}((\widetilde{E}_n/\widetilde{E}_{n-1} \wedge X)/C_p) \cong H_*(X).$$

In our case of interest, $X = \mathbb{T} \ltimes_{C_p} B^{\wedge p}$ is a $\mathbb{T}$ -equivariant spectrum, and we seek to understand the residual $\mathbb{T}/C_p$ -action on X^{tC_p} . The group $\mathbb{T}$ acts (only) on the even-indexed

terms in the Greenlees filtration, hence $\mathbb{T}/C_p$ acts (only) on the odd-indexed terms in the homotopy limit above. Restricting to this coarser tower of spectra, we can express the C_p -Tate construction as the homotopy limit

$$X^{tC_p} \simeq \operatorname{holim}_{n \rightarrow -\infty} X^{tC_p}[2n-1]$$

of $\mathbb{T}/C_p$ -equivariant spectra. There are $\mathbb{T}/C_p$ -equivariant homotopy (co-)fibre sequences

$$[\tilde{E}_{2n}/\tilde{E}_{2n-2} \wedge F(E\mathbb{T}_+, X)]^{C_p} \rightarrow X^{tC_p}[2n-1] \rightarrow X^{tC_p}[2n+1], \quad (4.6)$$

and equivalences

$$\begin{aligned} [\tilde{E}_{2n}/\tilde{E}_{2n-2} \wedge F(E\mathbb{T}_+, X)]^{C_p} &\simeq [\tilde{E}_{2n}/\tilde{E}_{2n-2} \wedge X]^{C_p} \\ &\simeq (\tilde{E}_{2n}/\tilde{E}_{2n-2} \wedge X)/C_p \cong \Sigma^{2n-1}\mathbb{T}_+ \wedge_{C_p} X, \end{aligned}$$

for all n , as $\tilde{E}_{2n}/\tilde{E}_{2n-2}$ is C_p -free.

The map κ is then realized as the homotopy limit of a tower of maps

$$\kappa[2n-1]: \mathbb{T}/C_p \ltimes (B^{\wedge p})^{tC_p}[2n-1] \rightarrow (\mathbb{T} \ltimes_{C_p} B^{\wedge p})^{tC_p}[2n-1]$$

induced by the C_p -equivariant inclusion $i_p: B^{\wedge p} \rightarrow \mathbb{T} \ltimes_{C_p} B^{\wedge p}$ and the $\mathbb{T}/C_p$ -action on the displayed target. Passing to homotopy fibres as in (4.6), we see that κ in Tate filtrations $(2n-1)$ and $2n$ is represented by the map

$$\bar{\kappa}: \mathbb{T}/C_p \ltimes (\Sigma^{2n-1}\mathbb{T}_+ \wedge_{C_p} B^{\wedge p}) \longrightarrow \Sigma^{2n-1}\mathbb{T}_+ \wedge_{C_p} (\mathbb{T} \ltimes_{C_p} B^{\wedge p})$$

induced by $\operatorname{id} \wedge_{C_p} i_p$ and the $\mathbb{T}/C_p$ -action. More explicitly, this is the map

$$\bar{\kappa} = \Sigma^{2n-1}(\xi \ltimes_{C_p} \operatorname{id})$$

induced by the C_p -equivariant homeomorphism

$$\xi: \mathbb{T}/C_p \times \mathbb{T} \xrightarrow{\cong} \mathbb{T} \times_{C_p} \mathbb{T}$$

that takes $([z], w)$ to $[zw, z]$. Here $z, w \in \mathbb{T}$ and square brackets indicate C_p -orbits.

A cellular approximation to ξ induces the C_p -equivariant chain isomorphism

$$\xi_*: C_*(\mathbb{T}/C_p) \otimes C_*(\mathbb{T}) \xrightarrow{\cong} C_*(\mathbb{T}) \otimes_{C_p} C_*(\mathbb{T})$$

given by

$$e_j \otimes e_k \longmapsto e_k \otimes e_j,$$

for $(j, k) = (0, 0)$, $(0, 1)$ or $(1, 1)$, whereas

$$e_1 \otimes e_0 \longmapsto e_0 \otimes e_1 + e_1 \otimes Te_0$$

in the case $(j, k) = (1, 0)$. This follows by combining the cellular model $e_0 \mapsto e_0$, $e_1 \mapsto e_0 \otimes e_1 + e_1 \otimes Te_0$ for the diagonal map $z \mapsto (z, z)$ with the cellular model $e_j \otimes e_k \mapsto e_{j+k}$ for $j + k \leq 1$, $e_1 \otimes e_1 \mapsto 0$ for the multiplication map $(z, w) \mapsto zw$.

Recall also that we made a choice of a chain equivalence $H_*(B) \simeq C_*(B)$, inducing a chain equivalence $H_*(B)^{\otimes p} \simeq C_*(B)^{\otimes p} \simeq C_*(B^{\wedge p})$ that is compatible with the C_p -actions. Combining these facts, we see that $\bar{\kappa}$ has a chain level model

$$\bar{\kappa}_*: C_*(\mathbb{T}/C_p) \otimes \Sigma^{2n-1}C_*(\mathbb{T}) \otimes_{C_p} H_*(B)^{\otimes p} \xrightarrow{\cong} \Sigma^{2n-1}C_*(\mathbb{T}) \otimes_{C_p} C_*(\mathbb{T}) \otimes_{C_p} H_*(B)^{\otimes p}$$

given by

$$e_j \otimes \Sigma^{2n-1}e_k \otimes x \longmapsto \Sigma^{2n-1}e_k \otimes e_j \otimes x, \quad (4.7)$$

for $(j, k) = (0, 0)$, $(0, 1)$ or $(1, 1)$, and

$$e_1 \otimes \Sigma^{2n-1}e_0 \otimes x \longmapsto (\Sigma^{2n-1}e_0 \otimes e_1 + \Sigma^{2n-1}e_1 \otimes Te_0) \otimes x \quad (4.8)$$

in the case $(j, k) = (1, 0)$, where $x \in H_*(B)^{\otimes p}$. Note that, as a consequence of linearity,

$$e_1 \otimes \Sigma^{2n-1} e_0 \otimes x - e_0 \otimes \Sigma^{2n-1} e_1 \otimes Tx \longmapsto \Sigma^{2n-1} e_0 \otimes e_1 \otimes x.$$

Now consider a class $u^i t^r \otimes \alpha^{\otimes p}$ in $H_*^c((B^{\wedge p})^{tC_p})$. Let $n = -r$ and $k = 1 - i$. Then $u^i t^r \otimes \alpha^{\otimes p}$ lies in Tate filtration $-(i + 2r) = 2n - 1 + k$, and is represented by the homology class of

$$\Sigma^{2n-1} e_k \otimes \alpha^{\otimes p}$$

in $H_*(\Sigma^{2n-1} \mathbb{T}_+ \wedge_{C_p} B^{\wedge p})$. Hence, $e_j \otimes u^i t^r \otimes \alpha^{\otimes p}$ is represented by the class of

$$e_j \otimes \Sigma^{2n-1} e_k \otimes \alpha^{\otimes p}$$

in $H_*(\mathbb{T}/C_p \ltimes (\Sigma^{2n-1} \mathbb{T}_+ \wedge_{C_p} B^{\wedge p}))$.

Its image under $\bar{\kappa}_*$ is given by the formulas (4.7) and (4.8), with $x = \alpha^{\otimes p}$, and hence equals the class of

$$\Sigma^{2n-1} e_k \otimes e_j \otimes \alpha^{\otimes p}$$

in $H_*(\Sigma^{2n-1} \mathbb{T}_+ \wedge_{C_p} (\mathbb{T} \ltimes_{C_p} B^{\wedge p}))$ for $(j, k) = (0, 0)$, $(0, 1)$ or $(1, 1)$, and the class of

$$(\Sigma^{2n-1} e_0 \otimes e_1 + \Sigma^{2n-1} e_1 \otimes e_0) \otimes \alpha^{\otimes p},$$

for $(j, k) = (1, 0)$, as $T(\alpha^{\otimes p}) = \alpha^{\otimes p}$. It also follows that

$$(e_1 \otimes u t^r - e_0 \otimes t^r) \otimes \alpha^{\otimes p}$$

is represented by

$$(e_1 \otimes \Sigma^{2n-1} e_0 - e_0 \otimes \Sigma^{2n-1} e_1) \otimes \alpha^{\otimes p},$$

which maps under $\bar{\kappa}_*$ to $\Sigma^{2n-1} e_0 \otimes e_1 \otimes \alpha^{\otimes p}$.

To find the Tate representative of $\kappa_*(e_j \otimes u^i t^r \otimes \alpha^{\otimes p})$, we now use the cofibre sequence (4.4) and the associated cofibre sequence

$$\Sigma^{2n-1}(\mathbb{T} \ltimes_{C_p} B^{\wedge p}) \rightarrow \Sigma^{2n-1} \mathbb{T}_+ \wedge_{C_p} (\mathbb{T} \ltimes_{C_p} B^{\wedge p}) \rightarrow \Sigma^{2n}(\mathbb{T} \ltimes_{C_p} B^{\wedge p})$$

coming from the Tate filtration of the middle term. It shows that $\Sigma^{2n-1} e_k \otimes e_j \otimes \alpha^{\otimes p}$ in

$$H_*(\Sigma^{2n-1} \mathbb{T}_+ \wedge_{C_p} (\mathbb{T} \ltimes_{C_p} B^{\wedge p}))$$

has Tate representative $u^i t^r \otimes e_j \otimes \alpha^{\otimes p}$, where $i = 1 - k$ and $r = -n$. Noting that the case $(i, j) = (1, 1)$ corresponds to the case $(j, k) = (1, 0)$, and chasing the formulas, we get the asserted result. $\square$

THEOREM 4.11. *The map*

$$\omega^t: \mathbb{T}/C_p \ltimes (B^{\wedge p})^{tC_p} \longrightarrow \mathrm{THH}(B)^{tC_p}$$

induces the complete $\mathcal{A}_$ -comodule homomorphism*

$$\omega_*^t: H_*(\mathbb{T}/C_p) \otimes H_*^c((B^{\wedge p})^{tC_p}) \longrightarrow H_*^c(\mathrm{THH}(B)^{tC_p})$$

given, modulo Tate filtration in the target, by

$$\begin{aligned} e_0 \otimes u^i t^r \otimes \alpha^{\otimes p} &\longmapsto u^i t^r \otimes \alpha^p, \\ e_1 \otimes t^r \otimes \alpha^{\otimes p} &\longmapsto t^r \otimes \alpha^{p-1} \wedge \alpha, \\ (e_1 \otimes u t^r - e_0 \otimes t^r) \otimes \alpha^{\otimes p} &\longmapsto u t^r \otimes \alpha^{p-1} \wedge \alpha, \end{aligned}$$

for $i \in \{0, 1\}$, $r \in \mathbb{Z}$.

Proof. The proof is clear from Proposition 4.9, Definition 3.10 and the naturality of the homological Tate spectral sequence with respect to the map $D \circ \omega_p: \mathbb{T} \ltimes_{C_p} B^{\wedge p} \rightarrow \mathrm{THH}(B)$. $\square$

5. Cyclotomic structure

Recall from diagram (2.1) that we are interested in the natural map

$$\hat{\Gamma}: [\widetilde{E\mathbb{T}} \wedge X]^{C_p} \rightarrow [\widetilde{E\mathbb{T}} \wedge F(E\mathbb{T}_+, X)]^{C_p} = X^{t_{C_p}},$$

for $X \cong \mathrm{THH}(B)$. We now study the source of this map. This allows us to construct the commutative diagram (2.4), which appears in Theorem 5.3.

DEFINITION 5.1. Let $G \subseteq \mathbb{T}$ be a closed subgroup containing C_p . For each good G -equivariant prespectrum Y , with spectrification $X = LY$, the geometric fixed point prespectrum $\Phi^{C_p}(Y)$ is the G/C_p -equivariant prespectrum with

$$\Phi^{C_p}(Y)(V^{C_p}) = Y(V)^{C_p},$$

and the *geometric fixed point* spectrum $\Phi^{C_p}(X)$ is its spectrification $\Phi^{C_p}(X) = L\Phi^{C_p}(Y)$; see [21, II.9.7] and [19, § 2.1].

There is a natural equivalence of G/C_p -spectra

$$\bar{s}: [\widetilde{E\mathbb{T}} \wedge X]^{C_p} \xrightarrow{\cong} \Phi^{C_p}(X);$$

see [21, II.9.8] and the proof of [19, Lemma 2.1]. We are concerned with the cases $G = C_p$, $Y = (B_{\mathrm{pre}}^{\wedge p})^\tau$ and $G = \mathbb{T}$, $Y = \mathrm{sd}_p \mathrm{thh}(B)^\tau$, corresponding to $X = B^{\wedge p}$ and $X = \mathrm{sd}_p \mathrm{THH}(B)$ respectively.

With notation as in Definition 3.1, there are natural isomorphisms and maps

$$\begin{aligned} \mathrm{sd}_p \mathrm{thh}(B; S^V)_k^{C_p} &= \mathrm{thh}(B; S^V)_{p(k+1)-1}^{C_p} \\ &= \left(\mathrm{hocolim}_{\vec{n} \in I^{p(k+1)}} \mathrm{Map}(S^{n_0} \wedge \dots \wedge S^{n_{p(k+1)-1}}, B_{n_0} \wedge \dots \wedge B_{n_{p(k+1)-1}} \wedge S^V) \right)^{C_p} \\ &\cong \mathrm{hocolim}_{\vec{n} \in I^{k+1}} \mathrm{Map}((S^{n_0} \wedge \dots \wedge S^{n_k})^{\wedge p}, (B_{n_0} \wedge \dots \wedge B_{n_k})^{\wedge p} \wedge S^V)^{C_p} \\ &\rightarrow \mathrm{hocolim}_{\vec{n} \in I^{k+1}} \mathrm{Map}(S^{n_0} \wedge \dots \wedge S^{n_k}, B_{n_0} \wedge \dots \wedge B_{n_k} \wedge S^{V^{C_p}}) = \mathrm{thh}(B; S^{V^{C_p}})_k \end{aligned}$$

induced from the identifications $S^{n_0} \wedge \dots \wedge S^{n_k} \cong ((S^{n_0} \wedge \dots \wedge S^{n_k})^{\wedge p})^{C_p}$ and $B_{n_0} \wedge \dots \wedge B_{n_k} \cong ((B_{n_0} \wedge \dots \wedge B_{n_k})^{\wedge p})^{C_p}$. These define a natural map of prespectra

$$r'_k: \Phi^{C_p}(\mathrm{sd}_p \mathrm{thh}(B)_k) \rightarrow \mathrm{thh}(B)_k,$$

for each $k \geq 0$, and likewise after applying the natural good thickening.

LEMMA 5.2. Suppose that B is connective. The spectrum maps

$$r'_0: \Phi^{C_p}(B^{\wedge p}) \xrightarrow{\cong} B^{\wedge 1}$$

(corresponding to the case $k = 0$) and

$$r': \Phi^{C_p}(\mathrm{sd}_p \mathrm{THH}(B)) \xrightarrow{\cong} \mathrm{THH}(B)$$

(obtained from the cases $k \geq 0$ by geometric realization) are natural equivalences.

Proof. The second case is proved in [19, Proposition 2.5], and the first case is part of their proof. In more detail, they prove that the connectivity of the map

$$\Omega^{V^{C_p}-W} r'_0(V^{C_p}): \Omega^{V^{C_p}-W} \mathrm{thh}(B; S^V)_{p-1}^{C_p} \longrightarrow \Omega^{V^{C_p}-W} \mathrm{thh}(B; S^{V^{C_p}})_0$$

grows to infinity with V for each fixed W . (The connectivity hypothesis is implicitly used at the bottom of p. 42 in the cited paper.) $\square$

In the case $G = \mathbb{T}$, let $\rho: \mathbb{T} \rightarrow \mathbb{T}/C_p$ be the p th root isomorphism of groups, with inverse $\rho^{-1}: \mathbb{T}/C_p \rightarrow \mathbb{T}$ taking $[z]$ to z^p . The cyclic structures on $\mathrm{sd}_p \mathrm{thh}(B)_\bullet$ and $\mathrm{thh}(B)_\bullet$ induce a $\mathbb{T}/C_p$ -equivariant structure on $\Phi^{C_p}(\mathrm{sd}_p \mathrm{THH}(B))$ and a $\mathbb{T}$ -equivariant structure on $\mathrm{THH}(B)$. These are compatible, in the sense that the cyclotomic structure equivalence r' is ρ^{-1} -equivariant; see [19, Definition 2.2].

The C_p -equivariant map $\eta_p: B^{\wedge p} \rightarrow \mathrm{sd}_p \mathrm{THH}(B)$ is induced from the inclusion of 0-simplices $B_{\mathrm{pre}}^{\wedge p} \rightarrow \mathrm{sd}_p \mathrm{thh}(B)_\bullet$, and hence is compatible with $\bar{s}$ by naturality, and with the maps r'_0 and r' by the construction of the latter via geometric realization. Hence, the outer part of the following diagram commutes:

$$\begin{array}{ccccc} B^{\wedge 1} & \xrightarrow{i} & \mathbb{T} \ltimes B^{\wedge 1} & \xrightarrow{\omega} & \mathrm{THH}(B) \\ \simeq \uparrow r'_0 & & \simeq \uparrow \rho^{-1} \ltimes r'_0 & & \simeq \uparrow r' \\ \Phi^{C_p}(B^{\wedge p}) & \longrightarrow & \mathbb{T}/C_p \ltimes \Phi^{C_p}(B^{\wedge p}) & \longrightarrow & \Phi^{C_p}(\mathrm{sd}_p \mathrm{THH}(B)) \\ \simeq \uparrow \bar{s} & & \simeq \uparrow \mathrm{id} \ltimes \bar{s} & & \simeq \uparrow \bar{s} \\ [\widetilde{E\mathbb{T}} \wedge B^{\wedge p}]^{C_p} & \longrightarrow & \mathbb{T}/C_p \ltimes [\widetilde{E\mathbb{T}} \wedge B^{\wedge p}]^{C_p} & \longrightarrow & [\widetilde{E\mathbb{T}} \wedge \mathrm{sd}_p \mathrm{THH}(B)]^{C_p}. \end{array} \quad (5.1)$$

The right-hand horizontal maps are the unique equivariant extensions, given by the $\mathbb{T}/C_p$ -action on $[\widetilde{E\mathbb{T}} \wedge \mathrm{sd}_p \mathrm{THH}(B)]^{C_p}$ and $\Phi^{C_p}(\mathrm{sd}_p \mathrm{THH}(B))$, and the $\mathbb{T}$ -action on $\mathrm{THH}(B)$ respectively. These equivariant extensions are compatible, via the identity $\mathrm{id}: \mathbb{T}/C_p \rightarrow \mathbb{T}/C_p$ and the isomorphism $\rho^{-1}: \mathbb{T}/C_p \rightarrow \mathbb{T}$, respectively, in view of the equivariance properties of $\bar{s}$ and r' discussed above. Hence, the whole diagram commutes.

THEOREM 5.3. *Let B be a connective symmetric ring spectrum. There is a natural commutative diagram in the stable homotopy category:*

$$\begin{array}{ccccc} & & \eta & & \\ & \curvearrowright & & \curvearrowleft & \\ B & \xrightarrow{i} & \mathbb{T} \ltimes B & \xrightarrow{\omega} & \mathrm{THH}(B) \\ \epsilon_B \downarrow & & \downarrow \rho \ltimes \epsilon_B & & \downarrow \gamma \\ (B^{\wedge p})^{tC_p} & \xrightarrow{i'} & \mathbb{T}/C_p \ltimes (B^{\wedge p})^{tC_p} & \xrightarrow{\omega^t} & \mathrm{THH}(B)^{tC_p} \\ & \curvearrowleft & & \curvearrowright & \\ & & \eta^t & & \end{array}$$

where ϵ_B is the composite

$$B \xleftarrow{\simeq} [\widetilde{E\mathbb{T}} \wedge B^{\wedge p}]^{C_p} \xrightarrow{\hat{\Gamma}} (B^{\wedge p})^{tC_p} = R_+(B),$$

and γ is the composite

$$\mathrm{THH}(B) \xleftarrow{\simeq} [\widetilde{E\mathbb{T}} \wedge \mathrm{sd}_p \mathrm{THH}(B)]^{C_p} \xrightarrow{\hat{\Gamma}} (\mathrm{sd}_p \mathrm{THH}(B))^{tC_p} \cong \mathrm{THH}(B)^{tC_p}.$$

Proof. By naturality of the map $\hat{\Gamma}$ with respect to the inclusion of 0-simplices $\eta_p: B^{\wedge p} \rightarrow \mathrm{sd}_p \mathrm{THH}(B)$, we get that the outer part of the following diagram commutes:

$$\begin{array}{ccccc}
 & & [\mathrm{id} \wedge \eta_p]^{C_p} & & \\
 & \searrow & & \nearrow & \\
 [\widetilde{E\mathbb{T}} \wedge B^{\wedge p}]^{C_p} & \longrightarrow & \mathbb{T}/C_p \ltimes [\widetilde{E\mathbb{T}} \wedge B^{\wedge p}]^{C_p} & \longrightarrow & [\widetilde{E\mathbb{T}} \wedge \mathrm{sd}_p \mathrm{THH}(B)]^{C_p} \\
 \downarrow \hat{\Gamma} & & \downarrow \mathrm{id} \ltimes \hat{\Gamma} & & \downarrow \hat{\Gamma} \\
 (B^{\wedge p})^{t_{C_p}} & \xrightarrow{i'} & \mathbb{T}/C_p \ltimes (B^{\wedge p})^{t_{C_p}} & \xrightarrow{\omega_p^{t_{C_p}} \circ \kappa} & (\mathrm{sd}_p \mathrm{THH}(B))^{t_{C_p}} \\
 & \searrow & & \nearrow & \\
 & & \eta_p^{t_{C_p}} & &
 \end{array} \quad (5.2)$$

The right-hand column is $\mathbb{T}/C_p$ -equivariant, and the right-hand square is constructed to be the unique $\mathbb{T}/C_p$ -equivariant extension of the outer part. Hence, the whole diagram commutes.

By combining the right-hand parts of diagrams (5.1), (5.2) and (4.2), we get the central part of the commutative diagram below:

$$\begin{array}{ccc}
 \mathbb{T} \ltimes B^{\wedge 1} & \xrightarrow{\omega} & \mathrm{THH}(B) \\
 \uparrow \simeq \rho^{-1} \ltimes r'_0 \bar{s} & & \uparrow \simeq r' \bar{s} \\
 \mathbb{T}/C_p \ltimes [\widetilde{E\mathbb{T}} \wedge B^{\wedge p}]^{C_p} & \longrightarrow & [\widetilde{E\mathbb{T}} \wedge \mathrm{sd}_p \mathrm{THH}(B)]^{C_p} \\
 \downarrow \mathrm{id} \ltimes \hat{\Gamma} & & \downarrow \hat{\Gamma} \\
 \mathbb{T}/C_p \ltimes (B^{\wedge p})^{t_{C_p}} & \longrightarrow & \mathrm{sd}_p \mathrm{THH}(B)^{t_{C_p}} \\
 \searrow \omega^t & & \downarrow \cong D^{t_{C_p}} \\
 & & \mathrm{THH}(B)^{t_{C_p}}
 \end{array}$$

$\rho \ltimes \epsilon_B$ (left curved arrow from $\mathbb{T} \ltimes B^{\wedge 1}$ to $\mathbb{T}/C_p \ltimes (B^{\wedge p})^{t_{C_p}}$)
 γ (right curved arrow from $\mathrm{THH}(B)$ to $\mathrm{THH}(B)^{t_{C_p}}$)

This proves the theorem. $\square$

6. The comparison map

In this section, we compute the effect of the comparison map $\gamma: \mathrm{THH}(B) \rightarrow \mathrm{THH}(B)^{t_{C_p}}$ for $B = \mathrm{MU}$ and BP . The main results are Theorems 6.4 and 6.9.

Let $H = H\mathbb{F}_p$ be the mod p Eilenberg–Mac Lane spectrum, realized as a commutative symmetric ring spectrum. Recall [26] the structure

$$\mathcal{A}_* = H_*(H) = P(\bar{\xi}_k \mid k \geq 1) \otimes E(\bar{\tau}_k \mid k \geq 0)$$

of the dual Steenrod algebra, where $|\bar{\xi}_k| = 2p^k - 2$ and $|\bar{\tau}_k| = 2p^k - 1$. (This assumes that p is odd, we leave the details for $p = 2$ to the reader.) The Hopf algebra coproduct is given by the formulas

$$\begin{aligned}
 \psi(\bar{\xi}_k) &= \sum_{i+j=k} \bar{\xi}_i \otimes \bar{\xi}_j^i, \\
 \psi(\bar{\tau}_k) &= 1 \otimes \bar{\tau}_k + \sum_{i+j=k} \bar{\tau}_i \otimes \bar{\xi}_j^i,
 \end{aligned}$$

where $\bar{\xi}_0 = 1$.

6.1. The case of complex cobordism

Let MU be the complex cobordism spectrum, realized as a commutative symmetric ring spectrum [24, IV.2]. Recall [1, pp. 75–77] the $\mathcal{A}_*$ -comodule algebra isomorphism

$$H_*(\mathrm{MU}) \cong P(\bar{\xi}_k \mid k \geq 1) \otimes P(m_\ell \mid \ell \neq p^k - 1),$$

where m_ℓ is an $\mathcal{A}_*$ -comodule primitive of degree 2ℓ , for each $\ell \geq 1$ not of the form $p^k - 1$. Note that $H_*(\mathrm{MU})$ is concentrated in even degrees.

DEFINITION 6.1. For $\ell = p^k - 1$, let $m_\ell = \bar{\xi}_k$, so that $H_*(\mathrm{MU}) \cong P(m_\ell \mid \ell \geq 1)$.

LEMMA 6.2. There is an $\mathcal{A}_*$ -comodule algebra isomorphism

$$H_*(\mathrm{THH}(\mathrm{MU})) \cong H_*(\mathrm{MU}) \otimes E(\sigma m_\ell \mid \ell \geq 1).$$

The classes σm_ℓ are $\mathcal{A}_*$ -comodule primitive, for all $\ell \geq 1$.

Proof. We use the (first quadrant) Bökstedt spectral sequence

$$E_{*,*}^2 = HH_*(H_*(B)) \Longrightarrow H_*(\mathrm{THH}(B))$$

arising from the skeleton filtration on $\mathrm{THH}(B)$, that is, the filtration induced from the simplicial structure on $\mathrm{thh}(B)$. See [3, § 5] for the tools used in this computation, including the facts that σ is a derivation and that it commutes with the Dyer–Lashof operations Q^i .

The E^2 -term for $B = \mathrm{MU}$ is

$$E_{*,*}^2 = H_*(\mathrm{MU}) \otimes E(\sigma m_\ell \mid \ell \geq 1).$$

We have $E_{*,*}^2 = E_{*,*}^\infty$, as all algebra generators lie in filtrations less than or equal to 1. There are no algebra extensions, as σm_ℓ is in an odd degree, and hence is an exterior class for p odd by graded commutativity, and for $p = 2$ by the Dyer–Lashof calculation $(\sigma m_\ell)^2 = Q^{2\ell+1}(\sigma m_\ell) = \sigma Q^{2\ell+1}(m_\ell) = 0$, as $Q^{2\ell+1}(m_\ell) \in H_{4\ell+1}(\mathrm{MU}) = 0$. The $\mathcal{A}_*$ -coaction $\nu: H_*(X) \rightarrow \mathcal{A}_* \otimes H_*(X)$ for $X = \mathrm{THH}(\mathrm{MU})$ is given by $\nu(\sigma m_\ell) = (1 \otimes \sigma)\nu(m_\ell) = (1 \otimes \sigma)(1 \otimes m_\ell) = 1 \otimes \sigma m_\ell$ for $\ell \neq p^k - 1$, whereas

$$\nu(\sigma \bar{\xi}_k) = (1 \otimes \sigma) \left(\sum_{i+j=k} \bar{\xi}_i \otimes \bar{\xi}_j^{p^i} \right) = \sum_{i+j=k} \bar{\xi}_i \otimes \sigma(\bar{\xi}_j^{p^i}) = 1 \otimes \sigma \bar{\xi}_k,$$

as $\sigma(\bar{\xi}_j^{p^i}) = 0$ for $i \geq 1$. □

The following was proved by a different method in [14, 6.4].

PROPOSITION 6.3. The homological Tate spectral sequence

$$\widehat{E}_{*,*}^2(\mathrm{THH}(\mathrm{MU})) = \widehat{H}^{-*}(C_p; H_*(\mathrm{THH}(\mathrm{MU}))) \Longrightarrow H_*^c(\mathrm{THH}(\mathrm{MU})^{tC_p})$$

collapses at the $\widehat{E}^3 = \widehat{E}^\infty$ -term, with

$$\widehat{E}_{*,*}^\infty = \widehat{H}^{-*}(C_p; \mathbb{F}_p) \otimes P(m_\ell^p \mid \ell \geq 1) \otimes E(m_\ell^{p-1} \sigma m_\ell \mid \ell \geq 1).$$

Proof. The $\widehat{E}^2$ -term is

$$\widehat{E}_{*,*}^2 = \widehat{H}^{-*}(C_p; \mathbb{F}_p) \otimes P(m_\ell \mid \ell \geq 1) \otimes E(\sigma m_\ell \mid \ell \geq 1),$$

where $\widehat{H}^{-*}(C_p; \mathbb{F}_p) = E(u) \otimes P(t, t^{-1})$ for p odd. The classes $u^i t^r$ are all infinite cycles, as recalled in Remark 4.3. The d^2 -differentials are given by the formula

$$d^2(u^i t^r \otimes \alpha) = u^i t^{r+1} \otimes \sigma \alpha;$$

see, for example, [14, 3.2]. The homology of $P(m_\ell) \otimes E(\sigma m_\ell)$ with respect to σ is $P(m_\ell^p) \otimes E(m_\ell^{p-1} \sigma m_\ell)$, and so, by the Künneth formula,

$$\widehat{E}_{*,*}^3 = \widehat{H}^{-*}(C_p; \mathbb{F}_p) \otimes P(m_\ell^p \mid \ell \geq 1) \otimes E(m_\ell^{p-1} \sigma m_\ell \mid \ell \geq 1).$$

By Theorem 4.11 and Corollary 3.13 the map

$$\omega^t: \mathbb{T}/C_p \times (B^{\wedge p})^{tC_p} \rightarrow \mathrm{THH}(B)^{tC_p},$$

for $B = \mathrm{MU}$ takes the classes $e_0 \otimes 1 \otimes m_\ell^{\otimes p}$ and $e_1 \otimes 1 \otimes m_\ell^{\otimes p}$ in $H_*(\mathbb{T}/C_p) \otimes H_*^c((B^{\wedge p})^{tC_p})$ to classes represented by $1 \otimes m_\ell^p$ and $1 \otimes m_\ell^{p-1} \sigma m_\ell$ in the Tate spectral sequence respectively. Hence, the $\widehat{E}^3$ -term is generated by infinite cycles, and there cannot be any further differentials. $\square$

THEOREM 6.4. *The map*

$$\gamma: \mathrm{THH}(\mathrm{MU}) \rightarrow \mathrm{THH}(\mathrm{MU})^{tC_p}$$

induces a complete $\mathcal{A}_$ -comodule algebra homomorphism*

$$\gamma_*: H_*(\mathrm{THH}(\mathrm{MU})) \longrightarrow H_*^c(\mathrm{THH}(\mathrm{MU})^{tC_p}),$$

mapping

$$\sigma m_\ell \longmapsto (-1)^{\ell t(p-1)\ell} \otimes m_\ell^{p-1} \sigma m_\ell$$

modulo Tate filtration in the target, for each $\ell \geq 1$.

Proof. We use the commutative diagram

$$\begin{array}{ccc} \mathbb{T} \ltimes \mathrm{MU} & \xrightarrow{\omega} & \mathrm{THH}(\mathrm{MU}) \\ \rho \ltimes \epsilon_{\mathrm{MU}} \downarrow & & \downarrow \gamma \\ \mathbb{T}/C_p \ltimes (\mathrm{MU}^{\wedge p})^{tC_p} & \xrightarrow{\omega^t} & \mathrm{THH}(\mathrm{MU})^{tC_p} \end{array}$$

from Theorem 5.3 for $B = \mathrm{MU}$. Let $[\mathbb{T}] \in H_1(\mathbb{T})$ be the fundamental class, which maps by ρ_* to the fundamental class $e_1 \in H_1(\mathbb{T}/C_p)$. Then $\omega_*([\mathbb{T}] \otimes \alpha) = \sigma \alpha$ for all $\alpha \in H_*(\mathrm{MU})$, so we can compute $\gamma_*(\sigma \alpha)$ as the image under ω_*^t of $e_1 \otimes (\epsilon_{\mathrm{MU}})_*(\alpha)$. We make separate calculations for the cases $\alpha = \xi_k$ (with $\ell = p^k - 1$) and $\alpha = m_\ell$ (with $\ell \neq p^k - 1$).

Recall from [23, 3.2.1] that the homomorphism

$$\epsilon_*: H_*(\mathrm{MU}) \rightarrow R_+(H_*(\mathrm{MU}))$$

is given by the formula

$$\epsilon_*(\alpha) = \sum_{r=0}^{\infty} t^{-(p-1)r} \otimes (-1)^r P_*^r(\alpha)$$

in the homological Singer construction $R_+(H_*(\mathrm{MU}))$, where P_*^r is the homology operation dual to the r th Steenrod power. (The terms involving $(\beta P^r)_*(\alpha)$ vanish in this case, as $H_*(\mathrm{MU})$ is concentrated in even degrees.)

By Lemma 6.5, we obtain

$$\begin{aligned}\epsilon_*(\bar{\xi}_k) &= \sum_{i=0}^k t^{-(p^i-1)} \otimes \bar{\xi}_{k-i}^{p^i} \\ &= 1 \otimes \bar{\xi}_k + t^{-(p-1)} \otimes \bar{\xi}_{k-1}^p + \dots + t^{-(p^k-1)} \otimes 1\end{aligned}$$

in $R_+(H_*(\mathrm{MU}))$ for each $k \geq 1$. To control the terms with $1 \leq i \leq k$, it will be convenient to compare with the ϵ_* -image of $\bar{\xi}_{k-1}^p$. By Lemma 6.6, we obtain

$$\epsilon_*(\bar{\xi}_{k-1}^p) = \sum_{i=0}^{k-1} t^{-p(p^i-1)} \otimes \bar{\xi}_{k-1-i}^{p^{i+1}} = \sum_{i=1}^k t^{-(p^i-p)} \otimes \bar{\xi}_{k-i}^{p^i}.$$

Hence,

$$\epsilon_*(\bar{\xi}_k) = 1 \otimes \bar{\xi}_k + t^{-(p-1)} \cdot \epsilon_*(\bar{\xi}_{k-1}^p) \quad (6.1)$$

in $R_+(H_*(\mathrm{MU}))$ for each $k \geq 1$. Here the multiplication by $t^{-(p-1)}$ refers to the $R_+(H_*(S)) = \widehat{H}^{-*}(C_p; \mathbb{F}_p)$ -module structure on $R_+(H_*(\mathrm{MU}))$, coming from the S -module structure on MU .

Next, recall the isomorphism $R_+(H_*(\mathrm{MU})) \cong H_*^c((\mathrm{MU}^{\wedge p})^{tC_p})$ of [23, 5.14], taking $t^r \otimes \alpha$ to a preferred class represented by

$$(-1)^\ell t^{r+(p-1)\ell} \otimes \alpha^{\otimes p}$$

in the Tate spectral sequence, where $|\alpha| = 2\ell$ is assumed to be even. (The coefficient is more complicated when $|\alpha|$ is odd.) By Lunøe–Nielsen and Rognes [23, 5.12], the map $\epsilon_{\mathrm{MU}}: \mathrm{MU} \rightarrow (\mathrm{MU}^{\wedge p})^{tC_p}$ induces the composite of $\epsilon_*: H_*(\mathrm{MU}) \rightarrow R_+(H_*(\mathrm{MU}))$ and this isomorphism. Hence, the identity (6.1) tells us that the difference

$$(\epsilon_{\mathrm{MU}})_*(\bar{\xi}_k) - t^{-(p-1)} \cdot (\epsilon_{\mathrm{MU}})_*(\bar{\xi}_{k-1}^p)$$

is the preferred class represented by

$$t^{(p-1)(p^k-1)} \otimes \bar{\xi}_k^{\otimes p}$$

in the Tate spectral sequence converging to $H_*^c((\mathrm{MU}^{\wedge p})^{tC_p})$.

We now chase the class $[\mathbb{T}] \otimes \bar{\xi}_{k-1}^p$ in $H_*(\mathbb{T} \times \mathrm{MU})$ around the commutative square above. Going to the right,

$$\omega_*([\mathbb{T}] \otimes \bar{\xi}_{k-1}^p) = \sigma(\bar{\xi}_{k-1}^p) = 0$$

in $H_*(\mathrm{THH}(\mathrm{MU}))$, as σ is a derivation. Hence, the image under γ_* is also 0, which implies that

$$(\omega^t)_*(e_1 \otimes (\epsilon_{\mathrm{MU}})_*(\bar{\xi}_{k-1}^p)) = 0.$$

It follows by the $\widehat{H}^{-*}(C_p; \mathbb{F}_p)$ -module structure that

$$(\omega^t)_*(e_1 \otimes t^{-(p-1)} \cdot (\epsilon_{\mathrm{MU}})_*(\bar{\xi}_{k-1}^p)) = 0.$$

Finally, we chase the class $[\mathbb{T}] \otimes \bar{\xi}_k$ around the square to see that $\gamma_*(\sigma \bar{\xi}_k)$ equals

$$(\omega^t)_*(e_1 \otimes (\epsilon_{\mathrm{MU}})_*(\bar{\xi}_k)) = (\omega^t)_*(e_1 \otimes t^{(p-1)(p^k-1)} \otimes \bar{\xi}_k^{\otimes p}),$$

which, by Theorem 4.11 and Corollary 3.13, is represented by

$$t^{(p-1)(p^k-1)} \otimes \bar{\xi}_k^{p-1} \sigma \bar{\xi}_k$$

in the Tate spectral sequence converging to $H_*^c(\mathrm{THH}(\mathrm{MU})^{tC_p})$. This proves the theorem for $m_\ell = \bar{\xi}_k$, with $\ell = p^k - 1$, as $(-1)^\ell \equiv +1 \pmod p$ in these cases.

The remaining cases, of m_ℓ with $\ell \neq p^k - 1$, are simpler. We have

$$\epsilon_*(m_\ell) = 1 \otimes m_\ell$$

in the homological Singer construction, as these classes m_ℓ are $\mathcal{A}_*$ -comodule primitives. Hence, $(\epsilon_{\text{MU}})_*(m_\ell)$ is the preferred class represented by

$$(-1)^{\ell} t^{(p-1)\ell} \otimes m_\ell^{\otimes p}$$

in the Tate spectral sequence converging to $H_*^c((\text{MU}^{\wedge p})^{tC_p})$. Chasing $[\mathbb{T}] \otimes m_\ell$ around the commutative diagram, we find that $\gamma_*(\sigma m_\ell)$ equals

$$\omega_*^t(e_1 \otimes (-1)^{\ell} t^{(p-1)\ell} \otimes m_\ell^{\otimes p}) = (-1)^{\ell} t^{(p-1)\ell} \otimes m_\ell^{p-1} \sigma m_\ell.$$

□

LEMMA 6.5.

$$(-1)^r P_*^r(\bar{\xi}_k) = \begin{cases} \bar{\xi}_{k-i}^{p^i} & \text{if } r = (p^i - 1)/(p - 1), \\ 0 & \text{otherwise.} \end{cases}$$

Proof. The Steenrod power P^r is dual to ξ_1^r in the Milnor basis $(\xi^I \tau^J)$ for $\mathcal{A}_*$, so each term of the form $\xi_1^r \otimes \alpha''$ in the coaction $\nu(\alpha)$ contributes a term α'' in $P_*^r(\alpha)$. From the recursive relation $0 = \sum_{i+j=k} \xi_i^{p^j} \cdot \bar{\xi}_j$ for $k \geq 1$, we see that $\bar{\xi}_i \equiv (-1)^i \xi_1^{(p^i-1)/(p-1)}$ for $i \geq 0$, modulo the ideal $J(0) \subset \mathcal{A}_*$ generated by the ξ_k with $k \geq 2$ and the τ_k with $k \geq 1$. Hence, $\nu(\bar{\xi}_k) = \sum_{i+j=k} \bar{\xi}_i \otimes \bar{\xi}_j^{p^i}$ is congruent to

$$\sum_{i=0}^k (-1)^i \xi_1^{(p^i-1)/(p-1)} \otimes \bar{\xi}_{k-i}^{p^i},$$

and contributes $(-1)^i \bar{\xi}_{k-i}^{p^i}$ to $P_*^r(\bar{\xi}_k)$ precisely if $r = (p^i - 1)/(p - 1)$. In this case $(-1)^i \equiv (-1)^r \pmod{p}$. □

LEMMA 6.6.

$$(-1)^r P_*^r(\bar{\xi}_{k-1}^p) = \begin{cases} \bar{\xi}_{k-1-i}^{p^{i+1}} & \text{if } r = p(p^i - 1)/(p - 1), \\ 0 & \text{otherwise.} \end{cases}$$

Proof. The coaction $\nu(\bar{\xi}_{k-1}^p) = \sum_{i+j=k-1} \bar{\xi}_i^p \otimes \bar{\xi}_j^{p^{i+1}}$ is congruent to

$$\sum_{i=0}^{k-1} (-1)^i \xi_1^{p(p^i-1)/(p-1)} \otimes \bar{\xi}_{k-1-i}^{p^{i+1}},$$

and contributes $(-1)^i \bar{\xi}_{k-1-i}^{p^{i+1}}$ to $P_*^r(\bar{\xi}_{k-1}^p)$ precisely if $r = p(p^i - 1)/(p - 1)$. □

6.2. The Brown–Peterson case

Let BP be the p -local Brown–Peterson spectrum, realized as an E_4 symmetric ring spectrum [7, 8]. We could avoid using the E_4 structure on BP by appealing to the symmetric ring spectrum map $\text{MU} \rightarrow \text{BP}$ of [6] and naturality, as in [14, 6.4], but this would make some arguments longer.

Recall [28, 4.1.12] the $\mathcal{A}_*$ -comodule algebra isomorphism

$$H_*(\text{BP}) \cong P(\bar{\xi}_k \mid k \geq 1).$$

Note that $H_*(\text{BP})$ is concentrated in even degrees.

LEMMA 6.7. *There is an $\mathcal{A}_*$ -comodule algebra isomorphism*

$$H_*(\mathrm{THH}(\mathrm{BP})) \cong H_*(\mathrm{BP}) \otimes E(\sigma \bar{\xi}_k \mid k \geq 1).$$

The classes $\sigma \bar{\xi}_k$ are $\mathcal{A}_*$ -comodule primitive, for all $k \geq 1$.

Proof. This is similar to the MU case; see [3, 5.12]. □

PROPOSITION 6.8. *The homological Tate spectral sequence*

$$\widehat{E}_{*,*}^2(\mathrm{THH}(\mathrm{BP})) = \widehat{H}^{-*}(C_p; H_*(\mathrm{THH}(\mathrm{BP}))) \implies H_*^c(\mathrm{THH}(\mathrm{BP})^{tC_p})$$

collapses at the $\widehat{E}^3 = \widehat{E}^\infty$ -term, with

$$\widehat{E}_{*,*}^\infty = \widehat{H}^{-*}(C_p; \mathbb{F}_p) \otimes P(\bar{\xi}_k^p \mid k \geq 1) \otimes E(\bar{\xi}_k^{p-1} \sigma \bar{\xi}_k \mid k \geq 1).$$

Proof. This is similar to the MU case; see also [14, 6.4]. □

THEOREM 6.9. *The map*

$$\gamma: \mathrm{THH}(\mathrm{BP}) \rightarrow \mathrm{THH}(\mathrm{BP})^{tC_p}$$

induces a complete $\mathcal{A}_*$ -comodule algebra homomorphism

$$\gamma_*: H_*(\mathrm{THH}(\mathrm{BP})) \longrightarrow H_*^c(\mathrm{THH}(\mathrm{BP})^{tC_p}),$$

mapping

$$\sigma \bar{\xi}_k \longmapsto t^{(p-1)(p^k-1)} \otimes \bar{\xi}_k^{p-1} \sigma \bar{\xi}_k,$$

modulo Tate filtration in the target, for each $k \geq 1$.

Proof. This is similar to the MU case, using the commutative diagram

$$\begin{array}{ccc} \mathbb{T} \ltimes \mathrm{BP} & \xrightarrow{\omega} & \mathrm{THH}(\mathrm{BP}) \\ \rho \ltimes \epsilon_{\mathrm{BP}} \downarrow & & \downarrow \gamma \\ \mathbb{T}/C_p \ltimes (\mathrm{BP}^{\wedge p})^{tC_p} & \xrightarrow{\omega^t} & \mathrm{THH}(\mathrm{BP})^{tC_p} \end{array}$$

from Theorem 5.3, or naturality with respect to the symmetric ring spectrum map $\mathrm{MU} \rightarrow \mathrm{BP}$. □

7. The Segal conjecture

In this section, we prove Theorem 2.2, which in turn implies Theorem 1.1. The main technical results are Propositions 7.2 and 7.3.

REMARK 7.1. The basic idea, in the case $B = \mathrm{MU}$, is that the homological Tate spectral sequences

$$' \widehat{E}_{*,*}^2 = \widehat{H}^{-*}(C_p; H_*(\mathrm{THH}(\mathrm{MU}))) \implies H_*^c(\mathrm{THH}(\mathrm{MU})^{tC_p})$$

and

$$'' \widehat{E}_{*,*}^2 = \widehat{H}^{-*}(C_p; H_*(\mathrm{THH}(\mathrm{MU})^{\wedge p})) \implies R_+(H_*(\mathrm{THH}(\mathrm{MU})))$$

have $\widehat{E}^\infty$ -terms

$$' \widehat{E}_{*,*}^\infty = E(u) \otimes P(t, t^{-1}) \otimes P(m_\ell^p \mid \ell \geq 1) \otimes E(m_\ell^{p-1} \sigma m_\ell \mid \ell \geq 1)$$

and

$$'' \widehat{E}_{*,*}^\infty = E(u) \otimes P(t, t^{-1}) \otimes P(m_\ell^{\otimes p} \mid \ell \geq 1) \otimes E(\sigma m_\ell^{\otimes p} \mid \ell \geq 1),$$

which are isomorphic by way of a filtration-shifting isomorphism given by

$$m_\ell^p \mapsto m_\ell^{\otimes p} \quad \text{and} \quad m_\ell^{p-1} \sigma m_\ell \mapsto t^m \otimes \sigma m_\ell^{\otimes p},$$

where $m = (p-1)/2$. The difficulty is to promote this isomorphism to an $\mathcal{A}_*$ -comodule isomorphism of the abutments, respecting the linear topologies. Our approach is to compare the two Tate towers via a third tower, given by base changing the Tate tower for $\mathrm{MU}^{\wedge p}$ along $\eta: \mathrm{MU} \rightarrow \mathrm{THH}(\mathrm{MU})$; see diagram (7.3).

Consider any connective symmetric ring spectrum B . We have a commutative diagram

$$\begin{array}{ccccc} \mathrm{THH}(B) & \xleftarrow{\eta} & B & \xrightarrow{\eta} & \mathrm{THH}(B) \\ \downarrow \epsilon_{\mathrm{THH}(B)} & & \downarrow \epsilon_B & & \downarrow \gamma \\ R_+(\mathrm{THH}(B)) & \xleftarrow{R_+(\eta)} & R_+(B) & \xrightarrow{\eta^t} & \mathrm{THH}(B)^{tC_p} \\ \downarrow & & \downarrow & & \downarrow \\ (\mathrm{THH}(B)^{\wedge p})^{tC_p}[n] & \xleftarrow{\quad} & (B^{\wedge p})^{tC_p}[n] & \xrightarrow{\quad} & \mathrm{THH}(B)^{tC_p}[n] \end{array} \quad (7.1)$$

by Theorem 5.3 and naturality of ϵ . We recall that $R_+(B) = (B^{\wedge p})^{tC_p}$, while $(B^{\wedge p})^{tC_p}[n]$ denotes the n th term in the Tate tower (4.5), and similarly with $\mathrm{THH}(B)$ in place of B . We are principally concerned with the limit behaviour as $n \rightarrow -\infty$.

Passing to continuous homology, we get a commutative diagram

$$\begin{array}{ccccc} H_*(\mathrm{THH}(B)) & \xleftarrow{\eta_*} & H_*(B) & \xrightarrow{\eta_*} & H_*(\mathrm{THH}(B)) \\ \downarrow \epsilon_* & & \downarrow \epsilon_* & & \downarrow \gamma_* \\ R_+(H_*(\mathrm{THH}(B))) & \xleftarrow{R_+(\eta_*)} & R_+(H_*(B)) & \xrightarrow{\eta_*^t} & H_*^c(\mathrm{THH}(B)^{tC_p}) \\ \downarrow & & \downarrow & & \downarrow \\ F^n R_+(H_*(\mathrm{THH}(B))) & \xleftarrow{\quad} & F^n R_+(H_*(B)) & \xrightarrow{\quad} & F^n H_*^c(\mathrm{THH}(B)^{tC_p}) \end{array} \quad (7.2)$$

of complete $\mathcal{A}_*$ -comodules. Here

$$F^n R_+(H_*(B)) = \mathrm{im}(H_*^c(R_+(B)) \rightarrow H_*((B^{\wedge p})^{tC_p}[n]))$$

so that $R_+(H_*(B)) = \lim_n F^n R_+(H_*(B))$, and similarly with $\mathrm{THH}(B)$ in place of B . We also use the notation

$$F^n H_*^c(\mathrm{THH}(B)^{tC_p}) = \mathrm{im}(H_*^c(\mathrm{THH}(B)^{tC_p}) \rightarrow H_*^c(\mathrm{THH}(B)^{tC_p}[n])),$$

so that $H_*^c(\mathrm{THH}(B)^{tC_p}) = \lim_n F^n H_*^c(\mathrm{THH}(B)^{tC_p})$.

Suppose now that B is an E_2 symmetric ring spectrum, so that $\mathrm{THH}(B)$ is a ring spectrum [13] and the upper two rows of (7.1) form a diagram of ring spectra. Then the upper two rows of (7.2) form a diagram of complete $\mathcal{A}_*$ -comodule algebras. Using the algebra structures, we

get a commutative diagram given by the solid arrows below:

$$\begin{array}{ccc}
 & H_*(\mathrm{THH}(B)) & \\
 \epsilon_* \swarrow & \downarrow & \searrow \gamma_* \\
 & R_+(H_*(B)) \otimes_{H_*(B)} H_*(\mathrm{THH}(B)) & \\
 f \swarrow & & \searrow g \\
 R_+(H_*(\mathrm{THH}(B))) & \xrightarrow[\Phi_B]{\quad} & H_*^c(\mathrm{THH}(B)^{tC_p}).
 \end{array} \tag{7.3}$$

Here $f(\alpha \otimes \beta) = R_+(\eta_*)(\alpha) \cdot \epsilon_*(\beta)$ while $g(\alpha \otimes \beta) = \eta_*^t(\alpha) \cdot \gamma_*(\beta)$. We wish to construct a suitably structured isomorphism $\Phi_B: R_+(H_*(\mathrm{THH}(B))) \rightarrow H_*^c(\mathrm{THH}(B)^{tC_p})$ making the whole diagram commute.

7.1. The case $B = \mathrm{MU}$

In this case the central term in diagram (7.3) is

$$R_+(H_*(\mathrm{MU})) \otimes_{H_*(\mathrm{MU})} H_*(\mathrm{THH}(\mathrm{MU})) \cong R_+(H_*(\mathrm{MU})) \otimes E(\sigma m_\ell \mid \ell \geq 1),$$

as $H_*(\mathrm{THH}(\mathrm{MU})) \cong H_*(\mathrm{MU}) \otimes E(\sigma m_\ell \mid \ell \geq 1)$.

By Lemma 6.2 each σm_ℓ is $\mathcal{A}_*$ -comodule primitive, and so, by Lunøe–Nielsen and Rognes [23, 3.2.1], we have $\epsilon_*(\sigma m_\ell) = 1 \otimes \sigma m_\ell$ in $R_+(H_*(\mathrm{THH}(\mathrm{MU})))$ on the left-hand side. It has Tate representative

$$t^{m(2\ell+1)} \otimes \sigma m_\ell^{\otimes p}$$

(up to a unit in $\mathbb{F}_p$) in Tate filtration $-(p-1)(2\ell+1)$, by Lunøe–Nielsen and Rognes [23, 5.14], where $m = (p-1)/2$. On the right-hand side, we showed in Theorem 6.4 that $\gamma_*(\sigma m_\ell)$ has Tate representative

$$t^{(p-1)\ell} \otimes m_\ell^{p-1} \sigma m_\ell$$

(up to a sign) in Tate filtration $-2(p-1)\ell$.

As both $\epsilon_*(\sigma m_\ell)$ and $\gamma_*(\sigma m_\ell)$ are in negative Tate filtration, we find that f maps $\alpha \otimes \beta$, with α in Tate filtration strictly less than n and $\beta \in E(\sigma m_\ell \mid \ell \geq 1)$, to a class in Tate filtration strictly less than n , and likewise for g . Hence, we get a commutative diagram:

$$\begin{array}{ccccc}
 R_+(H_*(\mathrm{THH}(\mathrm{MU}))) & \xleftarrow{f} & R_+(H_*(\mathrm{MU})) \otimes E(\sigma m_\ell \mid \ell \geq 1) & \xrightarrow{g} & H_*^c(\mathrm{THH}(\mathrm{MU})^{tC_p}) \\
 \downarrow & & \downarrow & & \downarrow \\
 F^n R_+(H_*(\mathrm{THH}(\mathrm{MU}))) & \xleftarrow{f_n} & F^n R_+(H_*(\mathrm{MU})) \otimes E(\sigma m_\ell \mid \ell \geq 1) & \xrightarrow{g_n} & F^n H_*^c(\mathrm{THH}(\mathrm{MU})^{tC_p}).
 \end{array}$$

Each f_n and g_n is an $\mathcal{A}_*$ -comodule homomorphism. Theorem 2.2 for MU now follows from the proposition below by letting

$$\Phi_{\mathrm{MU}} = \hat{g} \circ \hat{f}^{-1} = \left(\lim_n g_n \right) \circ \left(\lim_n f_n \right)^{-1}.$$

Recall the completed tensor product $\widehat{\otimes}$ from [23, § 2.5].

PROPOSITION 7.2. *The homomorphisms*

$$f_n: F^n R_+(H_*(\mathrm{MU})) \otimes E(\sigma m_\ell \mid \ell \geq 1) \longrightarrow F^n R_+(H_*(\mathrm{THH}(\mathrm{MU})))$$

and

$$g_n: F^n R_+(H_*(\mathrm{MU})) \otimes E(\sigma m_\ell \mid \ell \geq 1) \longrightarrow F^n H_*^c(\mathrm{THH}(\mathrm{MU})^{tC_p})$$

define strict maps $\{f_n\}_n$ and $\{g_n\}_n$ of inverse systems of bounded below $\mathcal{A}_*$ -comodules of finite type. These are pro-isomorphisms, in each homological degree. Hence, each of the limiting homomorphisms

$$\hat{f} = \lim_n f_n: R_+(H_*(\mathrm{MU})) \hat{\otimes} E(\sigma m_\ell \mid \ell \geq 1) \longrightarrow R_+(H_*(\mathrm{THH}(\mathrm{MU})))$$

and

$$\hat{g} = \lim_n g_n: R_+(H_*(\mathrm{MU})) \hat{\otimes} E(\sigma m_\ell \mid \ell \geq 1) \longrightarrow H_*^c(\mathrm{THH}(\mathrm{MU}))^{tC_p}$$

is a continuous isomorphism of complete $\mathcal{A}_*$ -comodules, with continuous inverse.

Proof. We begin with the proof for $\hat{f}$. The part of the Tate spectral sequence $\hat{E}^\infty$ -term

$$\begin{aligned} \hat{E}_{*,*}^\infty &= E(u) \otimes P(t, t^{-1}) \otimes P(m_\ell^{\otimes p} \mid \ell \geq 1) \\ &\implies R_+(H_*(\mathrm{MU})) \end{aligned}$$

in bidegrees (s, t) with Tate filtration $s \geq n$ is the associated graded of the preferred filtration of $F^n R_+(H_*(\mathrm{MU}))$. Hence, the part of

$$E(u) \otimes P(t, t^{-1}) \otimes P(m_\ell^{\otimes p} \mid \ell \geq 1) \otimes E(\sigma m_\ell \mid \ell \geq 1)$$

in Tate filtration $s \geq n$ is the associated graded of the filtration of $F^n R_+(H_*(\mathrm{MU})) \otimes E(\sigma m_\ell \mid \ell \geq 1)$. Similarly, the part of the Tate spectral sequence $\hat{E}^\infty$ -term

$$\begin{aligned} \hat{E}_{*,*}^\infty &= E(u) \otimes P(t, t^{-1}) \otimes P(m_\ell^{\otimes p} \mid \ell \geq 1) \otimes E(\sigma m_\ell^{\otimes p} \mid \ell \geq 1) \\ &\implies R_+(H_*(\mathrm{THH}(\mathrm{MU}))) \end{aligned}$$

in Tate filtration $s \geq n$ is the associated graded of the filtration of $F^n R_+(H_*(\mathrm{THH}(\mathrm{MU})))$.

The $\mathcal{A}_*$ -comodule homomorphism f_n is the identity on $F^n R_+(H_*(\mathrm{MU}))$, and takes σm_ℓ to $\epsilon_*(\sigma m_\ell)$ represented by a unit times $t^{m(2\ell+1)} \otimes \sigma m_\ell^{\otimes p}$ modulo filtration. Note that $t^{m(2\ell+1)} \otimes \sigma m_\ell^{\otimes p}$ lies in bidegree $(-(p-1)(2\ell+1), p(2\ell+1))$, on the line of slope $-p/(p-1)$ through the origin in the (s, t) -plane.

Now restrict attention to one total degree d , indicated by subscripts. We thus have compatible $\mathbb{F}_p$ -linear homomorphisms

$$f_{n,d}: [F^n R_+(H_*(\mathrm{MU})) \otimes E(\sigma m_\ell \mid \ell \geq 1)]_d \longrightarrow F^n R_+(H_*(\mathrm{THH}(\mathrm{MU})))_d,$$

for all integers n . To provide a pro-inverse, we shall define compatible $\mathbb{F}_p$ -linear homomorphisms

$$\phi_{n,d}: F^N R_+(H_*(\mathrm{THH}(\mathrm{MU})))_d \longrightarrow [F^n R_+(H_*(\mathrm{MU})) \otimes E(\sigma m_\ell \mid \ell \geq 1)]_d$$

with $N = N(n, d) = p(n-d) + d$ for all integers n . Write the source of the inclusion

$$[R_+(H_*(\mathrm{MU})) \otimes E(\epsilon_*(\sigma m_\ell) \mid \ell \geq 1)]_d \longrightarrow R_+(H_*(\mathrm{THH}(\mathrm{MU})))_d$$

as a direct sum

$$\bigoplus_L [R_+(H_*(\mathrm{MU})) \otimes \mathbb{F}_p\{\epsilon_L\}]_d$$

with L ranging over the strictly increasing sequences $L = (\ell_1 < \dots < \ell_r)$ of natural numbers, of length $r \geq 0$. Here

$$\epsilon_L = \epsilon_*(\sigma m_{\ell_1}) \cdot \dots \cdot \epsilon_*(\sigma m_{\ell_r})$$

has bidegree $(s_L, t_L) = (-(p-1)(2|L|+r), p(2|L|+r))$, where $|L| = \ell_1 + \dots + \ell_r$. The inclusion descends to an isomorphism

$$\bigoplus_L [F^{N-s_L} R_+(H_*(\mathrm{MU})) \otimes \mathbb{F}_p\{\epsilon_L\}]_d \xrightarrow{\cong} F^N R_+(H_*(\mathrm{THH}(\mathrm{MU})))_d,$$

as can be seen from the $\widehat{E}^\infty$ -terms. As $F^{N-s_L}R_+(H_*(\mathrm{MU}))$ is concentrated in total degrees greater than or equal to $N - s_L$, only the summands with $(N - s_L) + (s_L + t_L) \leq d$ are non-zero, and this is equivalent to $2|L| + r \leq d - n$. This inequality, in turn, implies that $N - s_L \leq n$. Hence, we have homomorphisms

$$\begin{aligned} [F^{N-s_L}R_+(H_*(\mathrm{MU})) \otimes \mathbb{F}_p\{\epsilon_L\}]_d &\hookrightarrow [F^{N-s_L}R_+(H_*(\mathrm{MU})) \otimes E(\sigma m_\ell \mid \ell \geq 1)]_d \\ &\twoheadrightarrow [F^n R_+(H_*(\mathrm{MU})) \otimes E(\sigma m_\ell \mid \ell \geq 1)]_d \end{aligned}$$

taking ϵ_L to $\sigma m_{\ell_1} \cdots \sigma m_{\ell_r}$. Taking the direct sum over L , and factoring through the isomorphism displayed above, we get the desired homomorphism $\phi_{n,d}$.

For varying n , the collection $\{\phi_{n,d}\}_n$ defines a pro-map, such that $f_{n,d} \circ \phi_{n,d}$ is equal to the structural surjection

$$F^N R_+(H_*(\mathrm{THH}(\mathrm{MU})))_d \twoheadrightarrow F^n R_+(H_*(\mathrm{THH}(\mathrm{MU})))_d,$$

and $\phi_{n,d} \circ f_{n,d}$ is equal to the structural surjection

$$[F^N R_+(H_*(\mathrm{MU})) \otimes E(\sigma m_\ell \mid \ell \geq 1)]_d \twoheadrightarrow [F^n R_+(H_*(\mathrm{MU})) \otimes E(\sigma m_\ell \mid \ell \geq 1)]_d.$$

Hence, $\{f_{n,d}\}_n$ is a pro-isomorphism, with pro-inverse $\{\phi_{n,d}\}_n$, in each total degree d .

The proof for $\hat{g}$ relies on similar filtration shift estimates. The associated graded of the filtration of $F^n R_+(H_*(\mathrm{MU})) \otimes E(\sigma m_\ell \mid \ell \geq 1)$ was discussed in the first part of the proof. The part of the Tate spectral sequence $\widehat{E}^\infty$ -term

$$\begin{aligned} \widehat{E}_{*,*}^\infty &= E(u) \otimes P(t, t^{-1}) \otimes P(m_\ell^p \mid \ell \geq 1) \otimes E(m_\ell^{p-1} \sigma m_\ell \mid \ell \geq 1) \\ &\implies H_*^c(\mathrm{THH}(\mathrm{MU})^{tC_p}) \end{aligned}$$

in Tate filtration $s \geq n$ is the associated graded of the filtration of $F^n H_*^c(\mathrm{THH}(\mathrm{MU})^{tC_p})$.

The $\mathcal{A}$ -comodule homomorphism g_n identifies $F^n R_+(H_*(\mathrm{MU}))$ with the Tate filtration $s \geq n$ part of $E(u) \otimes P(t, t^{-1}) \otimes P(m_\ell^p \mid \ell \geq 1)$, taking $m_\ell^{\otimes p}$ to m_ℓ^p . Furthermore, it takes σm_ℓ to $\gamma_*(\sigma m_\ell)$, represented by a sign times $t^{(p-1)\ell} \otimes m_\ell^{p-1} \sigma m_\ell$ modulo filtration. Here $t^{(p-1)\ell} \otimes m_\ell^{p-1} \sigma m_\ell$ lies in bidegree $(-2(p-1)\ell, 2p\ell + 1)$, on the line of slope $-p/(p-1)$ through the point $(s, t) = (0, 1)$.

In total degree d we have the strict pro-map $\{g_{n,d}\}_n$, with components

$$g_{n,d}: [F^n R_+(H_*(\mathrm{MU})) \otimes E(\sigma m_\ell \mid \ell \geq 1)]_d \longrightarrow F^n H_*^c(\mathrm{THH}(\mathrm{MU})^{tC_p})_d.$$

We define an $\mathbb{F}_p$ -linear pro-inverse $\{\psi_{n,d}\}_n$, with components

$$\psi_{n,d}: F^N H_*^c(\mathrm{THH}(\mathrm{MU})^{tC_p})_d \longrightarrow [F^n R_+(H_*(\mathrm{MU})) \otimes E(\sigma m_\ell \mid \ell \geq 1)]_d.$$

Here $N = N(n, d) = p(n - d) + d$, as in the $\hat{f}$ -case.

Write the source of the inclusion

$$[R_+(H_*(\mathrm{MU})) \otimes E(\gamma_*(\sigma m_\ell) \mid \ell \geq 1)]_d \longrightarrow H_*^c(\mathrm{THH}(\mathrm{MU})^{tC_p})_d$$

as a direct sum

$$\bigoplus_L [R_+(H_*(\mathrm{MU})) \otimes \mathbb{F}_p\{\gamma_L\}]_d$$

with $L = (\ell_1 < \dots < \ell_r)$ as above and

$$\gamma_L = \gamma_*(\sigma m_{\ell_1}) \cdots \gamma_*(\sigma m_{\ell_r})$$

in bidegree $(s'_L, t'_L) = (-2(p-1)|L|, 2p|L| + r)$. The inclusion descends to an isomorphism

$$\bigoplus_L [F^{N-s'_L} R_+(H_*(\mathrm{MU})) \otimes \mathbb{F}_p\{\gamma_L\}]_d \xrightarrow{\cong} F^N H_*^c(\mathrm{THH}(\mathrm{MU})^{tC_p})_d.$$

Only the summands with $(N - s'_L) + (s'_L + t'_L) \leq d$ are non-zero, and this implies $2|L| \leq d - n$ as $r \geq 0$. This, in turn, implies $N - s'_L \leq n$. Hence, we have homomorphisms

$$\begin{aligned} [F^{N-s'_L} R_+(H_*(\text{MU})) \otimes \mathbb{F}_p\{\gamma_L\}]_d &\hookrightarrow [F^{N-s'_L} R_+(H_*(\text{MU})) \otimes E(\sigma m_\ell \mid \ell \geq 1)]_d \\ &\twoheadrightarrow [F^n R_+(H_*(\text{MU})) \otimes E(\sigma m_\ell \mid \ell \geq 1)]_d \end{aligned}$$

taking γ_L to $\sigma m_{\ell_1} \cdots \sigma m_{\ell_r}$. Summing over L , and using the isomorphism above, we get the required homomorphism $\psi_{n,d}$.

The collection $\{\psi_{n,d}\}_n$ defines a pro-map, such that $g_{n,d} \circ \psi_{n,d}$ is equal to the structural surjection

$$F^N H_*^c(\text{THH}(\text{MU})^{tC_p})_d \twoheadrightarrow F^n H_*^c(\text{THH}(\text{MU})^{tC_p})_d,$$

and $\psi_{n,d} \circ g_{n,d}$ is equal to the structural surjection

$$[F^N R_+(H_*(\text{MU})) \otimes E(\sigma m_\ell \mid \ell \geq 1)]_d \twoheadrightarrow [F^n R_+(H_*(\text{MU})) \otimes E(\sigma m_\ell \mid \ell \geq 1)]_d.$$

Hence, $\{g_{n,d}\}_n$ is a pro-isomorphism. $\square$

7.2. The case $B = \text{BP}$

In this case, the central term in diagram (7.3) is

$$R_+(H_*(\text{BP})) \otimes_{H_*(\text{BP})} H_*(\text{THH}(\text{BP})) \cong R_+(H_*(\text{BP})) \otimes E(\sigma \bar{\xi}_k \mid k \geq 1).$$

By Lemma 6.7 each $\sigma \bar{\xi}_k$ is $\mathcal{A}_*$ -comodule primitive, so by Lunøe–Nielsen and Rognes [23, 3.2.1] we have $\epsilon_*(\sigma \bar{\xi}_k) = 1 \otimes \sigma \bar{\xi}_k$ in $R_+(H_*(\text{THH}(\text{BP})))$. It has Tate representative

$$t^{m(2p^k-1)} \otimes \sigma \bar{\xi}_k^{\otimes p}$$

in Tate filtration $-(p-1)(2p^k-1)$, by Lunøe–Nielsen and Rognes [23, 5.14]. We showed in Theorem 6.9 that $\gamma_*(\sigma \bar{\xi}_k)$ has Tate representative

$$t^{(p-1)(p^k-1)} \otimes \bar{\xi}_k^{p-1} \sigma \bar{\xi}_k$$

in Tate filtration $-2(p-1)(p^k-1)$. As both images are in negative Tate filtration, we get a commutative diagram

$$\begin{array}{ccccc} R_+(H_*(\text{THH}(\text{BP}))) & \xleftarrow{f} & R_+(H_*(\text{BP})) \otimes E(\sigma \bar{\xi}_k \mid k \geq 1) & \xrightarrow{g} & H_*^c(\text{THH}(\text{BP})^{tC_p}) \\ \downarrow & & \downarrow & & \downarrow \\ F^n R_+(H_*(\text{THH}(\text{BP}))) & \xleftarrow{f_n} & F^n R_+(H_*(\text{BP})) \otimes E(\sigma \bar{\xi}_k \mid k \geq 1) & \xrightarrow{g_n} & F^n H_*^c(\text{THH}(\text{BP})^{tC_p}). \end{array}$$

Theorem 2.2 for BP now follows from the proposition below by letting

$$\Phi_{\text{BP}} = \hat{g} \circ \hat{f}^{-1} = \left(\lim_n g_n \right) \circ \left(\lim_n f_n \right)^{-1}.$$

PROPOSITION 7.3. *The homomorphisms*

$$f_n: F^n R_+(H_*(\text{BP})) \otimes E(\sigma \bar{\xi}_k \mid k \geq 1) \longrightarrow F^n R_+(H_*(\text{THH}(\text{BP})))$$

and

$$g_n: F^n R_+(H_*(\text{BP})) \otimes E(\sigma \bar{\xi}_k \mid k \geq 1) \longrightarrow F^n H_*^c(\text{THH}(\text{BP})^{tC_p})$$

define strict maps $\{f_n\}_n$ and $\{g_n\}_n$ of inverse systems of bounded below $\mathcal{A}_*$ -comodules of finite type. These are pro-isomorphisms, in each homological degree. Hence, each of the limiting homomorphisms

$$\hat{f} = \lim_n f_n: R_+(H_*(BP)) \hat{\otimes} E(\sigma\bar{\xi}_k \mid k \geq 1) \longrightarrow R_+(H_*(\mathrm{THH}(BP)))$$

and

$$\hat{g} = \lim_n g_n: R_+(H_*(BP)) \hat{\otimes} E(\sigma\bar{\xi}_k \mid k \geq 1) \longrightarrow H_*^c(\mathrm{THH}(BP)^{tC_p})$$

is a continuous isomorphism of complete $\mathcal{A}_*$ -comodules, with continuous inverse.

Proof. The proof is similar to that of the MU case. □

References

1. J. F. ADAMS, *Stable homotopy and generalised homology*, Chicago Lectures in Mathematics (University of Chicago Press, Chicago, IL, 1974).
2. J. F. ADAMS, J. H. GUNAWARDENA and H. MILLER, ‘The Segal conjecture for elementary abelian p -groups’, *Topology* 24 (1985) 435–460.
3. V. ANGELTVEIT and J. ROGNES, ‘Hopf algebra structure on topological Hochschild homology’, *Algebr. Geom. Topol.* 5 (2005) 1223–1290.
4. CH. AUSONI and J. ROGNES, ‘Algebraic K -theory of topological K -theory’, *Acta Math.* 188 (2002) 1–39.
5. CH. AUSONI and J. ROGNES, ‘The chromatic red-shift in algebraic K -theory’, *Guido’s Book of Conjectures*, Monographie de L’Enseignement Mathématique 40 (2008) 13–15.
6. A. BAKER and A. JEANNERET, ‘Brave new Hopf algebroids and extensions of MU-algebras’, *Homology Homotopy Appl.* 4 (2002) 163–173.
7. M. BASTERRA and M. A. MANDELL, ‘Homology of E_n ring spectra and iterated THH’, *Algebr. Geom. Topol.* 11 (2011) 939–981.
8. M. BASTERRA and M. A. MANDELL, ‘The multiplication in BP ’, Preprint, 2011, arXiv:1101.0023.
9. M. BÖKSTEDT, ‘Topological Hochschild homology’, Bielefeld preprint (ca. 1986).
10. M. BÖKSTEDT, R. R. BRUNER, S. LUNØE-NIELSEN and J. ROGNES, ‘On cyclic fixed points of spectra’, Preprint, 2007, arXiv:0712.3476.
11. M. BÖKSTEDT, W. C. HSIANG and I. MADSEN, ‘The cyclotomic trace and algebraic K -theory of spaces’, *Invent. Math.* 111 (1993) 465–539.
12. M. BÖKSTEDT and I. MADSEN, ‘Topological cyclic homology of the integers’, *Astérisque* 226 (1994) 7–8, 57–143. K -theory (Strasbourg, 1992).
13. M. BRUN, Z. FIEDOROWICZ and R. M. VOGT, ‘On the multiplicative structure of topological Hochschild homology’, *Algebr. Geom. Topol.* 7 (2007) 1633–1650.
14. R. R. BRUNER and J. ROGNES, ‘Differentials in the homological homotopy fixed point spectral sequence’, *Algebr. Geom. Topol.* 5 (2005) 653–690.
15. G. CARLSSON, ‘Equivariant stable homotopy and Segal’s Burnside ring conjecture’, *Ann. of Math.* (2) 120 (1984) 189–224.
16. J. CARUSO, J. P. MAY and S. B. PRIDDY, ‘The Segal conjecture for elementary abelian p -groups. II. p -adic completion in equivariant cohomology’, *Topology* 26 (1987) 413–433.
17. J. P. C. GREENLEES, ‘Representing Tate cohomology of G -spaces’, *Proc. Edinb. Math. Soc.* (2) 30 (1987) 435–443.
18. J. P. C. GREENLEES and J. P. MAY, ‘Generalized Tate cohomology’, *Mem. Amer. Math. Soc.* 113 (1995) viii+178.
19. L. HESSELHOLT and I. MADSEN, ‘On the K -theory of finite algebras over Witt vectors of perfect fields’, *Topology* 36 (1997) 29–101.
20. M. HOVEY, B. SHIPLEY and J. SMITH, ‘Symmetric spectra’, *J. Amer. Math. Soc.* 13 (2000) 149–208.
21. L. G. LEWIS JR., J. P. MAY and M. STEINBERGER, *Equivariant stable homotopy theory*, Lecture Notes in Mathematics 1213 (Springer, Berlin, 1986). With contributions by J. E. McClure.
22. W. H. LIN, D. M. DAVIS, M. E. MAHOWALD and J. F. ADAMS, ‘Calculation of Lin’s Ext groups’, *Math. Proc. Cambridge Philos. Soc.* 87 (1980) 459–469.
23. S. LUNØE-NIELSEN and J. ROGNES, ‘The topological Singer construction’, Preprint, 2010, arXiv:1010.5633.
24. J. P. MAY, *E_∞ ring spaces and E_∞ ring spectra*, Lecture Notes in Mathematics 577 (Springer, Berlin, 1977). With contributions by Frank Quinn, Nigel Ray, and Jørgen Tornehave.
25. J. E. MCCLURE and R. E. STAFFELDT, ‘On the topological Hochschild homology of $\mathrm{bu}_+^{\mathrm{f}}$ ’, *Amer. J. Math.* 115 (1993) 1–45.
26. J. MILNOR, ‘The Steenrod algebra and its dual’, *Ann. of Math.* (2) 67 (1958) 150–171.

- 27. D. C. RAVENEL, 'The Segal conjecture for cyclic groups and its consequences', *Amer. J. Math.* 106 (1984) 415–446. With an appendix by Haynes R. Miller.
- 28. D. C. RAVENEL, *Complex cobordism and stable homotopy groups of spheres*, Pure and Applied Mathematics 121 (Academic Press, Orlando, FL, 1986).
- 29. J. ROGNES, 'Trace maps from the algebraic K -theory of the integers (after M. Bökstedt)', *J. Pure Appl. Algebra* 125 (1998) 277–286.

Sverre Lunøe-Nielsen and John Rognes

Department of Mathematics

University of Oslo

NO-0316 Oslo

Norway

sverreln@math.uio.no

rogn@math.uio.no